



0 0301 0013934 1





FRESH-WATER ALGÆ

OF THE

UNITED STATES;

(EXCLUSIVE OF THE DIATOMACEÆ)

COMPLEMENTAL TO

DESMIDS OF THE UNITED STATES;

VOLUME II.

PLATES.

2300 ILLUSTRATIONS,

COVERING ONE HUNDRED AND FIFTY-SEVEN COLORED PLATES.

BY THE REV. FRANCIS WOLLE,

Member of the American Society of Microscopists.

BETHLEHEM, PA.:
THE COMENIUS PRESS.

1887.


58085

EXPLANATIONS AND REFERENCES,

PLATE LIV.

Figures magnified 500 diameters.

	PAGE.
Fig. 1. TETMEMORUS GIGANTEUS,	34
“ 2. DOCIDIUM ARCHERII	25
“ 3. DOCIDIUM NODULOSUM,	26
“ 4. DOCIDIUM WOODII,	26
“ 5. DOCIDIUM BACULUM, Var. Floridense,	26
Figs. 6, 7. STAUROSTRUM VESICULATUM,	42
“ 8, 9 SPHAEROSOMA PULCHELLUM, Var. bambusioides,	22
Fig 10-13. EUASTRUM PECTINATUM, front, lateral and transverse views,	36
“ 14-16. HYALOTHECA DISSILIENS, Var. hians,	21




EXPLANATIONS AND REFERENCES,

PLATE LV.

Figures magnified 250 diameters.

		PAGE.
Figs. 1, 2.	CLOSTERIUM DIANÆ,	25
“ 3, 4.	CLOSTERIUM TURGIDUM,	24
Fig. 5-8.	CLOSTERIUM STRIOLATUM.	24
“ 9.	CLOSTERIUM DELPONTII,	25
“ 10.	CLOSTERIUM DIDYMOTICUM,	25
“ 11.	CLOSTERIUM SUBCOSTATUM,	24
“ 12.	CLOSTERIUM LUNULA, Var. striatum,	24
Figs. 13, 14.	CLOSTERIUM ENSIS.	23
“ 15, 16.	CLOSTERIUM PRELONGUM,	24
Fig. 17.	CLOSTERIUM BREBISSEI,	24
“ 18-20.	CLOSTERIUM LANCEOLATUM,	24
“ 21.	CLOSTERIUM JUNCIDUM Var. gracillima-loevissima.	23
“ 22.	CLOSTERIUM PRONUM,	25







EXPLANATIONS AND REFERENCES,

PLATE LVI.

Figures magnified 375-500 diameters.

		PAGE.
Figs. 1, 2.	MICRASTERIAS SPECIOSA,	38
Fig. 3.	MICRASTERIAS APICULATA	39
Figs. 4, 5.	MICRASTERIAS CRUX MELETENSE,	40
Fig. 6.	MICRASTERIAS RABENHORSTII, Var.,	40
Figs 7, 8.	CYLINDRICOYSTIS TUMIDUM,	23
Fig. 9.	XANTHIDIUM FASCICULATUM, Var. Subalpinum,	34
Figs. 10, 11.	XANTHIDIUM COLUMBIANUM,	34
Fig. 12.	CCYLINDROCYSTIS DE BARYI,	27
Figs. 13, 14.	XANTHIDIUM TORREYI,	35






EXPLANATIONS AND REFERENCES,

PLATE LVII.

Figures magnified 500 diameters.

	PAGE.
Figs. 1, 2. STAURASTRUM WOLLEANUM,	44
“ 3, 4. STAURASTRUM CORNUTUM,	43
“ 5, 6. STAURASTRUM BACILLARE,	41
“ 7, 8. STAURASTRUM MINNESOTENSE	43
“ 9, 10. STAURASTRUM PSEUDOCRENATUM,	42
“ 11-13. STAURASTRUM MINNEAPOLIENSE,	45
“ 14, 15. STAURASTRUM CALYXOIDES,	46
“ 16, 17. STAURASTRUM ERASUM	43
Fig. 18-20. COSMARIUM INFLATUM,	27
Figs. 21, 22. STAURASTRUM XIPHIDIOPHORUM,	44
Fig. 23-25. COSMARIUM NORDSTEDTII,	30
Figs. 26, 27. COSMARIUM SPHALEROSTICHUM,	31
“ 28, 29. COSMARIUM BRAUNII, forma MAJOR,	29
“ 30, 31. COSMARIUM SENDTNERIANUM,	29
Fig. 32. COSMARIUM PERFORATUM,	28
Figs. 33, 34. COSMARIUM LOBATUM,	28
“ 35, 36. COSMARIUM LOEVE,	30
Fig. 37. COSMARIUM CIRCULARE,	28








EXPLANATIONS AND REFERENCES,

PLATE LVIII.

Figures magnified 500 diameters.

	PAGE.
Fig. 1-3. EUASTRUM VENTRICOSUM, zygo-pore, lateral and end views of cell,	38
Figs. 4, 5. EUASTRUM CRASSUM, Var. SCROBICULATUM, front and lateral views,	38
Fig. 6-8. EUASTRUM MAGNIFICUM, front, lateral and end views,	37
“ 9-11. EUASTRUM PURUM, two front and one lateral view,	37
Figs. 12, 13. EUASTRUM CUNEATUM, front and lateral views,	37




EXPLANATIONS AND REFERENCES,

PLATE LIX.

Figures magnified 500 diameters.

	PAGE.
Fig. 1-3. STAURASTRUM WOLLEANUM, Var. KISSIMENSE,	44
Figs. 4, 5. STAURASTRUM TOHOPEKALIGENSE, front, lateral and end views,	45
“ 6, 7. MICRASTERIAS FURCATA, Var. SIMPLEX,	40
“ 8, 9. STAURASTRUM PARADOXUM, Var. OSCEOLENSE,	45
Fig. 10. MICRASTERIAS VERRUCOSA,	41
“ 11. SPHAEROSOMA MONILIFORME,	22
“ 12. SPHAEROSOMA PULCHRUM, Var. CONSTRICTUM,	22
Figs. 13, 14. BAMBUSINA GRACILESCENS, with zygospores,	21
Fig. 15. MICRASTERIAS RINGENS, Var. SERRULATA,	41




EXPLANATIONS AND REFERENCES,

PLATE LX.

Figures magnified 500 diameters.

	PAGE.
Figs. 1, 2. <i>COSMARIUM WOLLEANUM</i> , Var. <i>GRANULIFERUM</i> , front and lateral views,	29
Fig. 3-5. <i>COSMARIUM PARDALIS</i> , front, lateral and transverse views,	31
Figs. 6, 7. <i>COSMARIUM RHOMBUSOIDES</i> , front and transverse views,	28
“ 8, 9. <i>STAURASTRUM NOVE CAESAREÆ</i> , front and vertical views,	
“ 10, 11. <i>COSMARIUM QUINARIUM</i> front and vertical views.	32
Fig. 12-14. <i>COSMARIUM SUBCRUCIFORME</i> , front, lateral and vertical views,	33
Figs. 15, 16. <i>COSMARIUM AMERICANUM</i> , front and vertical views,	29
“ 17, 18. <i>COSMARIUM TAXICHONDRUM</i> , Var. <i>BIDENTULUM</i> , front and vertical views,	32
Fig. 19. <i>STAURASTRUM XIPHIOPHORUM</i> , Var. <i>SIMPLEX</i> , front view,	44
Figs. 20, 21. <i>COSMARIUM MICROSPHINCTUM</i> , front and vertical views,	32
“ 22, 23. <i>STAURASTRUM LUTEOLUM</i> , front and vertical views,	43
“ 24, 25. <i>COSMARIUM EXCAVATUM</i> , Var. <i>TRIGONUM</i> , front and vertical views,	30
“ 26, 27. <i>COSMARIUM LAGOENSE</i> , front and vertical views,	34
Fig. 28. <i>COSMARIUM CORDANUM</i> , front view,	27
“ 29. <i>EUASTRUM WOLLEI</i> , Var. <i>QUADRIGIBBERUM</i> , vertical view,	37
“ 30. <i>MICRASTERIAS AMERICANA</i> , Var. <i>SPINOSA</i> , front view of semi-cell,	39
“ 31-33. <i>COSMARIUM POLYMORPHUM</i> , front, lateral and vertical views,	31
Figs. 34, 35. <i>COSMARIUM OCTOGONUM</i> , front and vertical views,	33
Fig. 36-38. <i>STAURASTRUM ORBICULARE</i> , Var., oblique and front views,	41
Figs. 39, 40. <i>STAURASTRUM BRASILIENSE</i> , Var. <i>TRIQUETRUM</i> , front and vertical views,	46
“ 41, 42. <i>STAURASTRUM MUTICUM</i> , Var. <i>ELLIPTICUM</i> , front and vertical views.	41






EXPLANATIONS AND REFERENCES,

PLATE LXI.

Figures magnified 500 diameters except Closterium, only 250 diameters.

	PAGE.
Figs. 1, 2. CLOSTERIUM ACEROSUM, forma striolatum with single and twin zygospores.	24
Fig. 3. CLOSTERIUM LINEATUM, Var. costatum, with twin zygospores,	25
Figs. 4, 5. ARTHRODESMUS NOTOCHONDRUS, in front and vertical views,	35
Fig. 6. ARTHRODESMUS INCRASSATUS, Var. cycladatus,	35
Figs. 7, 8. COSMARIUM OCULIFERUM, front and lateral views,	30
Fig. 9-11. PENIUM CRUCIFERUM, in different stages of division	23
Figs. 12, 13. ARTHRODESMUS PACHYCERUS front and lateral views,	36
“ 14, 15. ARTHRODESMUS TRIANGULARIS, front and vertical views,	36
Fig. 16. DOCIDIUM GEORGICUM,	26
“ 17. PENIUM SPIROSTRIOLATUM, var.,	22
“ 19. “ “	23
“ 18. DOCIDIUM VERTICILLATUM,	27
Figs. 20, 21. DOCIDIUM RECTUM,	26






EXPLANATIONS AND REFERENCES,

PLATE LXII.

Figures magnified 500 diameters; the *Micrasterias* 375 diameters.


	PAGE.
Fig. 1. <i>MICRASTERIAS ALATA</i>	40
“ 2. <i>MICRASTERIAS MAMILLATA</i> ,	39
“ 3. <i>MICRASTERIAS DENTICULATA</i> , zygospore,	39
“ 4. <i>STAURASTRUM ARCTISCON</i> , var.,	45
“ 5. <i>COSMARIUM CONSPERSUM</i> , Var. <i>retusum</i> , front view,	31
“ 6. <i>COSMARIUM BIRETUM</i> , Var. <i>Floridense</i> , front view,	34
Figs. 7, 8. <i>STAURASTRUM COSMARIOIDES</i> , front and vertical views,	42
“ 9, 10. <i>STAURASTRUM BREVISPINA</i> , Var. <i>inermis</i> ,	42
“ 11, 12. <i>STAURASTRUM CORNUTUM</i> front and vertical views.	43



EXPLANATIONS AND REFERENCES,

PLATE LXIII.

		PAGE.
Fig. 1.	LEMANEA TORULOSA, natural size,	52
" 2.	" " magnified slightly,	52
" 3.	" " small portion of filament under higher power,	52
" 4.	" " portion of transverse section,	52
" 5.	LEMANEA FLUVIATILIS, slightly magnified,	52
" 6.	BATRACHOSPERMUM MONILIFORME, natural size,	57
" 10.	" " the same magnified.	
" 7.	BATRACHOSPERMUM, Var. pulcherrimum, natural size,	57
" 8.	" " " small section magnified,	57
" 9.	BATRACHOSPERMUM, Var. atrum, somewhat magnified.	57







EXPLANATIONS AND REFERENCES,

PLATE LXIV.

		PAGE.
Fig. 1.	BATRACHOSPERMUM VAGUM, Var. keratophytum, natural size,	58
“ 2.	“ “ end of a filament magnified, .	58
“ 3.	“ “ two whorls of same under higher power,	58
Figs. 4, 5.	Var. KUEHNIANUM, sections moderately magnified, .	57
Fig. 6.	BATRACHOSPERMUM MONILIFORME, Var. atrum, natural size. The same magnified, Plate LXIII, fig. 9,	57







EXPLANATIONS AND REFERENCES,

PLATE LXV.

	PAGE.
Figs 1, 2. BATRACHOSPERMUM VAGUM, developed stems,	57
Fig. 3-6. " " younger filaments,	57







EXPLANATIONS AND REFERENCES,

PLATE LXVI.

		PAGE.
Fig. 1.	TUOMEYA FLUVIATILIS,	53
“ 2.	TUOMEYA GRANDE, nearly twice the natural size,	54
“ 3.	“ “ small part of a filament enlarged; the outer membrane removed to show the in- ternal structure,	54
Figs. 4-6.	“ “ supporting branchlets at the nodes,	54
“ 7, 8.	“ “ part of a cord drawn out of the filament,	54






EXPLANATIONS AND REFERENCES,

PLATE LXVII.


		PAGE.
Fig. 1.	CHANTRANSIA PYGMAEA, natural size,	61
Figs. 2, 3.	“ “ two filaments,	61
“ 4, 5.	“ “ clusters of carpogons,	61
Fig. 6.	CHANTRANSIA VIRGATULA, magnified 125 diameters,	61
“ 7.	“ “ tetrasporangia	61
“ 8.	“ “ natural size, growing on a marine plant,	61
“ 9-11.	BANGIA ATRO-PURPUREA, four specimens in different stages of growth and cell division,	55




EXPLANATIONS AND REFERENCES,

PLATE LXVIII.

	PAGE.
Figs. 1, 2. CHANTRANSIA VIOLACEA, natural size,	59
Fig. 3. " " three filaments magnified about 125 diameters	59
Figs. 4, 5. Clusters of carpogons or sporangia emitting gonidia.	59
Fig. 6. Five single gonidia settled in a scratch on glass of aquarium and multiplied by cell division,	59
" 7. A cluster of gonidia under higher power, developing young plants,	59
" 8. A group of young plants,	59
Figs. 9, 10. Specimens of Var. C. expansa,	59
" 11, 12. Clusters of carpogons,	60
Fig. 13. Natural size of plants,	60
" 14. Var. C. Beardslei, magnified 125 diameters,	60






EXPLANATIONS AND REFERENCES,

PLATE LXIX.

	PAGE.
Fig. 1-7. CHANTRANSIA MACROSPORA, three filaments,	59
“ 2-4. Carpospores fertilized,	59
“ 5-7. Carpogons with trichogyne,	59
“ 8. Natural size,	59
“ 9. Section of old stem corticated,	59
“ 10-12. Matured carpospores in early stage of development,	59
“ 13. CHANTRANSIA HERMANNI, natural size,	60
“ 14. “ “ two filaments,	60
Figs. 15, 16. “ “ clusters of carpogons,	60
Fig. 17. HILDEBRANDTIA RIVULARIS, thallus peeled from river-stone, highly magnified,	62
Fig. 18. “ “ small transverse section under higher magnification, showing the spore receptacles,	62






EXPLANATIONS AND REFERENCES,

PLATE LXX.

	PAGE.
Fig. 1. COMPSOPOGON COERULEUS, a slightly magnified branched filament,	62
“ 2. End of a branch magnified 125 diameters,	62
“ 3. A section from lower part of filament, magnified 125 diameters,	62
“ 4. An older part of stem showing the corticulation, magnified 125 diameters,	62
“ 5. A section near the base, corticated, magnified 125 diameters,	62








EXPLANATIONS AND REFERENCES,

PLATE LXXI.


	PAGE.
Fig. 1. THOREA RAMOSISSIMA, slightly magnified,	58
Figs. 2, 3. Sections of filaments highly magnified,	58
“ 4, 5. Base of ramuli under greater magnification,	58




EXPLANATIONS AND REFERENCES,

PLATE LXXII.

	PAGE.
Figs. 1, 2. COLEOCHAETE SOLUTA, two sections of fronds,	64
Fig. 3. " " a loose growth in fruit,	64
Figs. 4, 5. " " two carpogons,	64
" 6, 7. " " two young forms developing from carpogons,	64
Fig. 8-13. COLEOCHAETE SCUTATA, younger and older plants of various forms,	64
" 14. COLEOCHAETE ORBICULARIS,	64
" 15-16. COLEOCHAETE IRREGULARIS,	65






EXPLANATIONS AND REFERENCES,

PLATE LXXIII.

	PAGE.
Fig. 1.	OEOGONIUM, filament of a nannandrous alga, 67
" 2.	Transverse striae the result of the process of cell multiplication, 67
" 3.	The same more fully illustrated, 67
" 4.	Filament of female nannandrous alga, 67
" 5.	Androsporangia with androspores escaping; oogonium with pore superior, 67
" 6.	Filament of female monœcious plant, 67
" 7.	Spermogonia hypogynous, spore spherical, pore superior, . . 67
" 8.	Filament of a monœcious form; 9. spermogonia cells hypogynous; 10. subhypogynous; 11. epigynous; 12. subepigynous; 13. apical. 67
" 14.	Pore of oogonium open centrally, 67
" 15.	A male plant, 67
" 16.	Androsporangia cells, 67
" 17.	Androspores escaping, 67
" 18-19.	Dwarf males. basal supporting cells and spermogonia, or upper cells of the dwarf males which bear the spermatozoids. opening on the top to allow their escape to fertilize the oospore, 67
" 20.	A zoospore making its way out of cell, 67
" 21-23.	Three zoospores fully out of cells, 67
" 24-26.	Zoospores in first stages of development, 67
" 27-30.	More advanced stages of growth, 67
" 31-33.	Imperfect zoospores passed out through slit in cell, remaining attached, 67
" 34.	An oogonium and oospore fertilized by a multitude of active spermatozoides (?) 67
" 35.	An oogonium showing how some are destitute of a pore, but open by an operculum or lid, 67
Figs. 36, 37.	An oospore passing out of an oogonium of a vegetative plant, 67
" 38, 39.	Young developments from oospores, 67







EXPLANATIONS AND REFERENCES,

PLATE LXXIV.

Figures magnified 250 diameters.

	PAGE
Figs. 1, 2. OEDOGONIUM CRYPTOPORUM, Var. <i>vulgare</i> ,	70
Fig. 3. OEDOGONIUM OBSOLETUM,	71
“ 4-6. OEDOGONIUM FRAGILE,	71
Figs. 7, 8. OEDOGONIUM ZIG-ZAG,	71
“ 9, 10. OEDOGONIUM VERNALE,	71
Fig. 11-14. OEDOGONIUM CRISPUM, Var. <i>rostellatum</i> ,	72
“ 15. “ “ type form,	72
“ 16-19. OEDOGONIUM POLYMORPHUM,	73
Figs. 20, 21. OEDOGONIUM PLUSIOSPORUM,	72








EXPLANATIONS AND REFERENCES,

PLATE LXXV.

Figures magnified 250 diameters.

	PAGE.
Fig. 1. OEDOGONIUM PALUDOSUM,	74
“ 2. OEDOGONIUM GRACILLIMUM,	74
“ 3. OEDOGONIUM LONGATUM,	95
“ 4-6. OEDOGONIUM FONTICOLA,	93
Figs. 7, 8. OEDOGONIUM LONDINENSE,	94
“ 9, 10. OEDOGONIUM PACHYANDRIUM,	89
Fig. 11-13. OEDOGONIUM CAPILLIFORME,	88






EXPLANATIONS AND REFERENCES,

PLATE LXXVI.

Figures magnified 250 diameters.

	PAGE.
Fig. 1. OEDOGONIUM GRANDE, oospore and two imperfect zoospores attached to filament,	95
Figs. 2, 3. OEDOGONIUM CRASSUM,	74
Fig. 4-6. OEDOGONIUM GIGANTEUM,	94
Figs. 7, 8. OEDOGONIUM RIVULARE, Var. major,	92




EXPLANATIONS AND REFERENCES.

PLATE LXXVII.

Figures magnified 250 diameters.

		PAGE.
Figs. 1, 2.	OEDOGONIUM PLATYGYNUM, two sections of filaments,	75
“ 3, 4.	“ “ transverse sections of oogonia,	75
“ 5, 6.	OEDOGONIUM DECIPIENS,	76
Fig. 7.	OEDOGONIUM ARESCHOUGH,	76
“ 8.	OEDOGONIUM UNDULATUM,	76
Figs. 9, 10.	OEDOGONIUM LUNDENSE,	79
Fig. 11-13.	OEDOGONIUM PYRIFORME,	95
“ 14-16.	OEDOGONIUM CRASSIUSCULUM,	80
“ 17.	“ “ an oospore surrounded by spermatozoides, active in the space between the spore and the membrane of the oogonium,	80
Figs. 18, 19.	Sections of filaments of male plants constituting the idioandrosporous form of the species,	80






EXPLANATIONS AND REFERENCES,

PLATE LXXVIII.

Figures magnified 250 diameters.

	PAGE.
Figs. 1, 2. OEDOGONIUM FLAVESCENS,	78
Fig. 3. OEDOGONIUM PRINCEPS,	93
Figs. 4, 5. OEDOGONIUM IRREGULARE,	79
Fig. 6-9. OEDOGONIUM BORISIANUM, in several forms and conditions,	81





EXPLANATIONS AND REFERENCES,

PLATE LXXIX.

Figures magnified 250 diameters.

	PAGE.
Figs. 1-3. OEDOGONIUM CONCATENATUM,	81
“ 4, 5. “ “ Var. setigerum,	82
“ 6, 7. OEDOGONIUM BRAUNII,	79
“ 8, 9. OEDOGONIUM SEXANGULARE,	82
“ 10, 11. OEDOGONIUM ACROSPORUM, Var. boreale,	84
Fig. 12-14. OEDOGONIUM OBTRUNCATUM,	84






EXPLANATIONS AND REFERENCES,

PLATE LXXX.

Figures magnified 250 diameters.

	PAGE.
Fig. 1-5. OEDOGONIUM WOLLEANUM, larger and smaller forms,	82
Figs. 6, 7. OEDOGONIUM MULTISPORA,	78




EXPLANATIONS AND REFERENCES,

PLATE LXXXI.

Figures magnified 250 diameters.

	PAGE.
Fig. 1-3. OEDOGONIUM AUTUMNALE, section of filaments,	73
Figs. 4, 5. " " first stages of development from zoospores,	73
" 6, 7. OEDOGONIUM OBSOLETUM, compare Plate LXXIV, fig. 3,	71
Fig. 8-11. OEDOGONIUM LANDBOROUGHII,	91
Figs. 12, 13. OEDOGONIUM DELICATULUM,	93
" 14, 15. OEDOGONIUM MONILIFORME,	93
" 16, 17. OEDOGONIUM RUFESCENS,	89
" 18, 19. OEDOGONIUM HEXAGONUM,	94
Fig. 20-22. OEDOGONIUM CYATHIGERUM,	77




EXPLANATIONS AND REFERENCES,


PLATE LXXXII.

Figures magnified 250 diameters

	PAGE.
Fig. 1-3. OEDOGONIUM MACRANDRIUM,	80
“ 4-6. OEDOGONIUM PRINGSHEIMII,	90
“ 7-9. OEDOGONIUM FRANKLINIANUM,	89
“ 10. OEDOGONIUM URBICUM,	74
“ 11-13. OEDOGONIUM BOSCHII,	91
Figs. 14, 15. OEDOGONIUM CARBONICUM,	90









EXPLANATIONS AND REFERENCES,

PLATE LXXXIII.

Figures magnified 250 diameters.

	PAGE.
Figs. 1, 2. OEDOGONIUM RIVULARE,	92
Fig. 3-6. OEDOGONIUM STAGNALE,	88
Figs. 7, 8. OEDOGONIUM CAPILLARE,	87
Fig. 9. OEDOGONIUM STAGNALE, Var.,	92






EXPLANATIONS AND REFERENCES,

PLATE LXXXIV.

Figures magnified 250 diameters.

	PAGE.
Figs. 1, 2. OEDOGONIUM STELLATUM,	85
Fig. 3-6. OEDOGONIUM DONNELLI,	85
“ 7. OEDOGONIUM ECHINOSPERMUM, type-form,	86
“ 8. OEDOGONIUM HYSTRIX	87
“ 9. OEDOGONIUM HUNTII,	85
“ 10 OEDOGONIUM CILIATUM,	84






EXPLANATIONS AND REFERENCES,

PLATE LXXXV.

Figures magnified 250 diameters.

	PAGE.
Figs. 1, 2. OEDOGONIUM ACROSPORUM, Var. Floridense,	83
Fig. 3-5. OEDOGONIUM PUNCTATO-STRIATUM,	91
“ 6-9. OEDOGONIUM ECHINOSPERMUM, Var. spermagonio-multicellulari,	86
“ 10-12. OEDOGONIUM CATARACTUM,	77
Figs. 13, 14. OEDOGONIUM ECHINATUM,	86





EXPLANATIONS AND REFERENCES,

PLATE LXXXVI.

Figures magnified 250 diameters.

	PAGE.
Fig. 1-3. BULBOCHAETE INTERMEDIA,	97
“ 4 BULBOCHAETE CRENULATA,	97
“ 5. BULBOCHAETE ELACHISTANDRA,	97
Figs. 6, 7. BULBOCHAETE SETIGERA,	98
Fig. 8-11. “ “ zoospores in different stages of exit,	






EXPLANATIONS AND REFERENCES,

PLATE LXXXVII.

Figures magnified 250 diameters.

		PAGE
Fig. 1.	BULBOCHAETE GIGANTEA,	99
Figs. 2, 3.	BULBOCHAETE MIRABILIS,	100
“ 4, 5.	BULBOCHAETE PYGMAEA,	100
Fig. 6.	BULBOCHAETE NANA,	100
“ 7.	BULBOCHAETE MINOR,	101









EXPLANATIONS AND REFERENCES,

PLATE LXXXVIII.

Figures magnified 250 diameters.

	PAGE.
Fig. 1. BULBOCHAETE BREBISSEONII,	99
Figs. 2, 3. BULBOCHAETE INSIGNIS,	101






EXPLANATIONS AND REFERENCES,


PLATE LXXXIX.

Figures magnified 250 diameters.

	PAGE.
Fig. 1-3. <i>BULBOCHAETE SETIGERA</i> , Var. <i>Canbyii</i> ,	99
“ 4-5. <i>BULBOCHAETE RHADINOSPORA</i> ,	103
“ 6-9. <i>BULBOCHAETE POLYANDRA</i> ,	98








EXPLANATIONS AND REFERENCES,

PLATE XC.

Figures magnified 250 diameters.

	PAGE.
Fig. 1-3. BULBOCHAETE RECTANGULARIS,	102
“ 4. BULBOCHAETE REPANDA,	102
“ 5. BULBOCHAETE SUBSIMPLEX,	101
Figs. 6, 7. BULBOCHAETE RECTANGULARIS, dwarf males, with spermagonia five-celled,	102






EXPLANATIONS AND REFERENCES,

PLATE XCI.

	PAGE.
Fig. 1-4. <i>CYLINDROCAPSA GEMINELLA</i> , cells of filaments developing	
oospores,	105
Figs. 5-6. Maturing oospores,	105
Fig. 7. Oospore escaping from tegument.	105
Figs. 8, 9. Supposed spermagonia,	105
Fig. 10-12. Oospores dividing and developing young forms,	105
Figs. 13, 14. Spore divided into two and four, the same enlarging previous to	
redivision,	105
Fig. 15-17. Redivided and growing,	105
“ 18-20. <i>CYLINDROCAPSA AMENA</i> , vegetative filaments and spore forma-	
tion	105
“ 21-23. Spore development,	105
“ 24. <i>PRASIOLA MEXICANA</i> , natural size,	107
“ 25. <i>PRASIOLA CRISPA</i> , natural size,	106
Figs. 26, 27. “ “ small sections magnified about 400 diameters, 106	







EXPLANATIONS AND REFERENCES,

PLATE XCII.

	PAGE.
Fig. 1. DRAPARNALDIA GLOMERATA, a stem with branches,	108
Figs. 2, 3. A part of a rhizome,	108
Fig. 4-6. Younger growths,	108
Figs. 7, 8. Stems producing zoospores,	108
Fig. 9-12. Ciliated zoospores and progressive stages of growth,	108






EXPLANATIONS AND REFERENCES,

PLATE XCIII.

	PAGE.
Fig. 1-4. DRAPARNALDIA SPINOSA, larger and smaller stems,	109
“ 5-8. “ “ branchlets of various sizes,	109
“ 3. A provision to hold larger branches to a stem by small bands,	109





EXPLANATIONS AND REFERENCES.

PLATE XCIV.


	PAGE.
Figs. 1, 2, (1-6). DRAPARNALDIA FLUMOSA, two robust specimens, .	. 109




EXPLANATIONS AND REFERENCES,

PLATE XCV.

	PAGE.
Fig. 1-4. DRAPARNALDIA RAVENELII, larger stems, more or less clothed with fascicles of branchlets. . .	110






EXPLANATIONS AND REFERENCES,

PLATE XCVI.

	PAGE.
Fig. 1. STIGEOCLONIUM THERMALE, a well-developed plant,	111
Figs. 2, 3. Microzoosporangia,	111
“ 4, 5. Microspores escaping and enlarging,	111
Fig. 6. A large sporangium,	111
“ 7-9. Early growth of Stigeoclonium,	111
“ 10. STIGEOCLONIUM NANUM,	112
“ 11. STIGEOCLONIUM TENUE,	111
“ 12. “ “ Var. <i>bulbiferum</i> ,	111
“ 13-16. The bulbous cells,	111






EXPLANATIONS AND REFERENCES,

PLATE XCVII.


	PAGE.
Figs. 1, 2. STIGEOCLONIUM FLAGELLIFERUM, type-form; and variety <i>crassiusculum</i> ,	112



EXPLANATIONS AND REFERENCES,

PLATE XCVIII.


	PAGE.
Figs. 1, 2, STIGEOCLONIUM NUDIUSCULUM. full length of plants, . . .	113
Fig. 3. " " end of branched filament with matured zoospore passed out of cells,	113
" 4. STIGEOCLONIUM AMGENUM,	113



EXPLANATIONS AND REFERENCES,
PLATE XCIX.

Figures magnified 250 diameters.

	PAGE.
Fig. 1. STIGEOCLONIUM FASCICULARE, a well-developed plant,	. 114
“ 2. STIGEOCLONIUM SUBSECUNDUM, full length plant, .	112




EXPLANATIONS AND REFERENCES,

PLATE C.


Reid

Figures magnified 250 diameters.

	PAGE.
Fig. 1. STIGEOCLONIUM FASTIGIATA. 114
Figs. 2, 3. STIGEOCLONIUM LONGIPILUS, according to Rabenhorst. Com- pare Plate CII. 115









EXPLANATIONS AND REFERENCES,


PLATE CI.

Figures magnified 250 diameters.

	PAGE.
Figs. 1, 2. STIGEOCLONIUM PROTENSUM,	112
“ 3, 4. “ “ the terminal setae,	112
Fig. 5-7. STIGEOCLONIUM TENUE, Var. <i>irregularis</i> ,	112
Figs. 8, 9. STIGEOCLONIUM UNIFORME, Var. <i>gracile</i> ,	111
Fig. 10-11. “ “ articulations developing sporangia,	111








EXPLANATIONS AND REFERENCES,

PLATE CII.

Figures magnified 250 diameters.

	PAGE.
Fig. 1-3. STIGEOCLONIUM LONGIPILUS, a form implied in Kirchner's diagnosis	114
" 4. STIGEOCLONIUM RADIAN,	115
" 5-8. CHAETOPHORA, developing long filaments,	114, 115







EXPLANATIONS AND REFERENCES,

PLATE CIII.

		PAGE.
Fig. 1-3.	CHAETOPHORA PISIFORMIS, thalli natural size, as frequently found attached to submerged grasses,	116, 114
Figs. 4, 5.	CHAETOPHORA ELEGANS, two sections of thalli largely magnified. Showing the arrangement of the filaments and character of branching,	116
Fig. 6.	A branch producing zoospores,	116
“ 7.	Natural size of thalli attached to a submerged leaf,	116
“ 8-10.	Three threads developing long, branched Stigeoclonium-like filaments. For higher developments or maturer growth, see Plate C, fig. 1,	116
“ 11.	CHAETOPHORA TUBERCULOSA, thalli of larger size,	116
“ 12-15.	CHAETOPHORA PISIFORMIS, thalli as they sometimes appear on the surface of river pebbles, primarily very small then increasing in size,	116
“ 16.	CHAETOPHORA LONGIPILA, natural size of thalli,	118
“ 17.	A thallus greatly magnified,	118
“ 18.	CHAETOPHORA MONILIFERA, a thallus magnified,	118
“ 19.	Thalli natural size,	118






EXPLANATIONS AND REFERENCES,

PLATE CIV.


			PAGE
Fig. 1.	CHAETOPHORA ENDIVIAEFOLIA,	Var. <i>cornuta</i> or <i>cornu damae</i> ,	
		twice the natural size,	117
“ 2.	“	“ a simple lobe of the same	
		greatly magnified,	117
“ 3.	“	“ Var. <i>linearis</i> ,	117
“ 4.	“	“ small section magnified	117
“ 5.	“	“ Var. <i>ramosissima</i> , a small form,	
		natural size,	117
Figs. 6, 7.	“	“ larger form, natural size,	117
“ 8, 9.	“	“ small section of a lobe, greatly	
		enlarged,	117




EXPLANATIONS AND REFERENCES,

PLATE CVI.

	PAGE.
Fig. 1-6. GONGROSIRA, a <i>pseudo</i> -genus,	1:0
Figs. 2, 3. GONGROSIRA SCLEROCOCCUS,	120
“ 7, 8. PROTONEMA, a prothallus of moss,	120
“ 10. “ “ of fern,	120







EXPLANATIONS AND REFERENCES,

PLATE CVII.

Figures magnified 125 diameters.

		PAGE.
Figs 1, 2.	CLADOPHORA FRACTA,	124
" 3, 4.	" " escaping zoogonidia,	124
" 5, 6.	" forma <i>gossypina</i>	125
" 7, 8.	CLADOPHORA FLOTOWIANA,	126








EXPLANATIONS AND REFERENCES,

PLATE CVII.

Figures magnified 125 diameters.

		PAGE.
Figs 1, 2.	CLADOPHORA FRACTA,	124
" 3, 4.	" " escaping zoogonidia,	124
" 5, 6.	" forma <i>gossypina</i>	125
" 7, 8.	CLADOPHORA FLOTOWIANA,	126





EXPLANATIONS AND REFERENCES,

PLATE CIX.

Figures magnified 125 diameters.

	PAGE.
Fig. 1-3. CLADOPHORA FRACTA, forma <i>viadrina</i> ,	125
“ 4-6. CLADOPHORA CRISPATA, forma <i>vitrea</i> ,	126
“ 7-10. “ “ “ <i>brachyclados</i> ,	126







EXPLANATIONS AND REFERENCES,

PLATE CX.

Figures magnified 125 diameters.

		PAGE.
Figs. 1, 2.	CLADOPHORA GLOMERATA,	. 128
Fig. 3.	“ forma <i>pumila</i> ,	. 128
“ 4-8.	“ peculiar development,	. 128






EXPLANATIONS AND REFERENCES,

PLATE CXI.

Figures magnified 125 diameters.

		PAGE.
Figs. 1, 2.	CLADOPHORA CANALICULARIS,	126
“ 3, 4.	CLADOPHORA GLOMERATA, forma <i>clavata</i> ,	128
Fig. 5.	“ “ “ <i>rivularis</i> ,	127




EXPLANATIONS AND REFERENCES,

PLATE CXII.

Figures magnified 125 diameters.

	PAGE.
Fig. 1-4. CLADOPHORA GLOMERATA, forma <i>simplicior</i> ,	127
“ 5. CLADOPHORA AEGAGROPILA,	129






EXPLANATIONS AND REFERENCES,

CXIII.

Figures magnified 125 diameters.

			PAGE.	
Fig. 1-3.	PITHOPHORA	OEDOGONIA, Var. <i>vaucherioides</i> with numerous		
		sporangia,	130	
Figs. 4, 5.	“	“	sporal branchlets,	131
“ 6, 7.	“	“	two immature filaments,	131




EXPLANATIONS AND REFERENCES,

PLATE CXIV.

Figures magnified 125 diameters.

		PAGE.
Fig. 1-5.	PITHOPHORA AEQUALIS, Var. <i>Floridense</i> . Sections of filaments with sporangia,	131
Figs. 6, 7.	PITHOPHORA KEWENSIS sterile filaments from Florida,	131
“ 8, 9.	“ “ section of fertile filament from Witrock's monograph,	131






EXPLANATIONS AND REFERENCES,

PLATE CXV.

Figures magnified 125 diameters.

	PAGE.
Fig. 1-4. CHROOLEPUS AUREUM, fruiting plants,	121
“ 5-7. Another form of the same species,	121
“ 8-15. Zoosporangia,	121
Figs. 16, 17. Zoospores escaping,	121
Fig. 18-21. Zoospores in different stages of copulation,	121
“ 22-25. C. AUREUM, Var. <i>corticulatum</i> ,	121
“ 26-29. Sporangial cells,	121
“ 30-33. C. MONILIFORME, Kg.,	123





EXPLANATIONS AND REFERENCES,

PLATE CXVI.

Figures magnified 125 diameters.

	PAGE.
Fig. 1-3. CHIROOLEPUS UMBRINUM, Kg., form frequent in Southern States,	123
“ 4-5. Pennsylvania form,	123
“ 6. C. ODORATUS,	122
Figs. 7, 8. C. LICHENICOLUS,	122
Fig. 9-12. C. IOLITHUS, Linn., <i>Veilchen-moss</i> ,	122







EXPLANATIONS AND REFERENCES,

PLATE CXVII.

Figures magnified 125 diameters.

		PAGE.
Fig. 1-3.	ULOTHRIX ZONATA, three vegetative filaments,	133
“ 4.	Cells of filament with macrozoospores,	134
“ 5.	Three zoospores,	134
“ 6-9.	Zoospores in early stage of development,	134
“ 10.	Cells of filament with microzoospores	134
“ 11.	The same enclosed in cysts,	134
“ 12.	Separated and at rest,	134
Figs. 13, 14.	Microzoospores germinating in cells,	134
“ 15, 16.	Zygospores,	134
Fig. 17-19.	Microzoospores conjugating—two of them fertilized,	134
“ 20-22.	ULOTHRIX YOUNGANUM,	135
“ 23-25.	ULOTHRIX SPECIOSA,	135
“ 26.	Partly empty cells from which the macrozoospores passed out,	135







EXPLANATIONS AND REFERENCES,


PLATE CXVIII.

Figures magnified 125 diameters.

	PAGE.
Figs. 1, 2. ULOTHRIX TENUIS,	134
Fig. 3-5. ULOTHRIX AEQUALIS,	134
“ 6-8. ULOTHRIX RIVULARIS,	136
Figs. 9, 10. ULOTHRIX SUBTILIS,	136
“ 11, 12. ULOTHRIX SUBTILISSIMA,	136
“ 13, 14. ULOTHRIX, Var. <i>stagnorum</i> ,	136
“ 15, 16. ULOTHRIX, Var. <i>variabilis</i> ,	136
Fig. 17. ULOTHRIX, Var. <i>tenerrima</i> ,	136
“ 18-19. ULOTHRIX THERMARUM,	136
“ 20-21. ULOTHRIX COMPACTA,	136
“ 22-26. ULOTHRIX SUBTILIS, an unusual form,	136
Figs. 27, 28. ULOTHRIX FLACCIDA—U. SUBTILIS,	137, 135
Fig. 29-33. ULOTHRIX RIVULARIS, Var. <i>cataracta</i> ,	137
“ 34-36. ULOTHRIX OSCILLARINA,	137









EXPLANATIONS AND REFERENCES,

PLATE CXIX.

Figures magnified 125 diameters.

	PAGE.
Figs. 1-2. ULOTHRIX NITENS,	137
Fig. 3-7. ULOTHRIX VARIA,	138
“ 8-10. ULOTHRIX PARIETINA, young forms,	138
“ 11-15. “ “ older forms,	138
“ 16-18. SCHIZOGONIUM MURALE,	137
“ 19-20. ULOTHRIX LENORMANDI,	138






EXPLANATIONS AND REFERENCES,

PLATE CXX.

Figures magnified 125 diameters.

	PAGE.
Figs. 1, 2. CONFERVA TENERRIMA,	143
“ 3, 4. CONFERVA RHYPOPHILA,	143
Fig. 5-8. CONFERVA GLACIALIODES,	143
Figs. 9, 10. CONFERVA FUNKII,	142
Fig. 11-13. CONFERVA AFFINIS,	141
“ 14-16. CONFERVA UTRICULOSA,	140
“ 17-20. CONFERVA FONTINALIS,	141
“ 21-24. CONFERVA FLOCCOSA, usual condition prior to division,	140
Figs. 23, 25. “ “ after division,	140
“ 26, 27. CONFERVA, joints forming into sporangial cells	140
“ 29, 30. CONFERVA, filaments in a state of dissolution,	140
Fig. 31-33. CONFERVA FUGACISSIMA,	141








EXPLANATIONS AND REFERENCES,

PLATE CXXI.

Figures magnified 125 diameters.

	PAGE.
Fig. 1-3. CONFERVA AMENA, normal vegetative condition,	140
“ 4. Older filament in state of dissolution; cells separating, microgonidia being scattered,	140
“ 5. Microgonidia under high power,	140
Figs. 6, 7. CONFERVA VULGARIS, the type-form,	142
“ 8, 9. CONFERVA BOMBYCINA,	142
“ 10, 11. CONFERVA VULGARIS, Var. <i>Farlowii</i> ,	142
Fig. 12. “ “ microgonidia escaping,	142
“ 13. “ “ young forms,	142
“ 14-17. “ “ stages of decay— <i>Hormospora</i> forms,	142
Figs. 18, 19. CONFERVA ABBREVIATA,	143
“ 20, 21. CONFERVA PUNCTALIS,	142
Fig. 22-25. RHIZOCLONIUM FONTINALE.	144
“ 26-28. RHIZOCLONIUM FLUITANS,	145
Figs. 29, 30. RHIZOCLONIUM HIEROGLYPHICUM,	144
“ 31, 32. “ “ Var. <i>Americana</i> ,	144






EXPLANATIONS AND REFERENCES,

PLATE CXXII.

Figures magnified 125 diameters.

	PAGE.
Fig. 1-4. RHIZOCLONIUM MAJOR, 146
“ 5-7. RHIZOCLONIUM SALINUM, 145
“ 8-10. RHIZOCLONIUM STAGNALE, 145
“ 11-12. RHIZOCLONIUM CASPARYI, 145
“ 13-16. RHIZOCLONIUM HOSFORDII, 145






EXPLANATIONS AND REFERENCES,

PLATE CXXIII.

Figures magnified 125 diameters.

	PAGE.
Fig. 1. SPHAEROPLEA ANNULINA, filament chlorophyllous green,	104
“ 2. Shows the arrangement of the mature (red) spores,	104
“ 3. Frequent appearance of filament preceding development of spermatozooids,	104
“ 4. Spermatozooids set free, more highly magnified,	104
“ 5. Zoogonidia under high power,	104
“ 6-9. APIOCYSTIS BRAUNIANA, two young fronds; two mature fronds, one of them with cells undergoing segmentation and a large frond with matured gonidia,	202
“ 10. Free ciliated gonidia or zoospores,	202
Figs. 11, 12. UROCOCCUS INSIGNIS,	201
Fig. 13. UROCOCCUS HOOKERIANUS,	201





EXPLANATIONS AND REFERENCES,

PLATE CXXIV.

Figures magnified 250 diameters.

	PAGE.
Fig. 1-4. BOTRYCOCCUS BRAUNII, smaller and larger clusters, . . .	195
“ 5-7. BOTRYDINA VULGARIS, younger and older thalli, . . .	195
Figs. 8, 9. PALMODACTYLON VARIUM,	189
Fig. 10-23. HORMOSPORA forms	(137, 133), 189





EXPLANATIONS AND REFERENCES,

PLATE CXXV.

Figures magnified 125 diameters.

	PAGE.
Fig. 1-6. SCHIZOMERIS LEIBLEINII, parts of filaments. The vegetative plant and broken ends with zoogonidia passing out	See below.
Figs. 7, 8. Young plants,
Fig. 9. ENTEROMORPHA INTESTINALIS	(308), 107
" 10. ENTEROMORPHA, small portion of membrane of older plant highly magnified, 107
" 11. CHAETOMORPHA, natural size, 143
" 12-13. Sections of filaments enlarged, 144
" 14. Escaping microgonidia, 144

[A description of the following genus was inadvertently omitted in the proper place, (page 17),].

Genus, SCHIZOMERIS, Kg.

Thallus chlorophyl green, filiform, composed primarily of a single series of cells which are usually broader than long, here and there strongly contracted; at first attached by a disc-like expansion of a narrow base, later often floating in mass; grows in the beginning by the division of the cells transversely, later in two or three directions.

Zoogonidia are formed in the cells of younger plants; they escape singly from the thinner filaments and in larger numbers from the larger ones; they are furnished with two or three cilia.


SCHIZOMERIS LEIBLEINII, Kg.

Bright green to deep blackish-green.

Diameter of filaments, 20-125 μ .

*Frequent in ditches and ponds, Pennsylvania, New Jersey, Florida, and probably in every State.

It is easily recognized by a *Ulothrix*-like look, which soon becomes familiar to the eye; it is larger than most of the species of *Ulothrix*, more irregular in outline by constrictions, the basal parts thin and the apices usually tapering to a point; the fruiting is also distinct.






EXPLANATIONS AND REFERENCES,


PLATE CXXXVI.

Figures magnified 166 diameters.

	PAGE.
Fig. 1-3. VAUCHERIA DICHOTOMA,	149
“ 4-6. “ “ sessile oogonia,	149
“ 7. “ “ escaping microgonidia,	149
“ 8-10. VAUCHERIA DILLWYNI,	150
“ 11-15. VAUCHERIA THURETHI,	149









EXPLANATIONS AND REFERENCES,
PLATE CXXVII.

Figures magnified 166 diameters.

	PAGE.
Fig. 1-4. VAUCHERIA PILUS, relative sizes of filaments and macrospores,	153
“ 5-8. VAUCHERIA AVERSA, with oogonia and antheridia,	150
“ 9-11. VAUCHERIA SESSILIS, three type-forms with oogonia and antheridia,	151
Figs. 12, 13. VAUCHERIA SERICEA, section of filament abundantly fruiting,	150







EXPLANATIONS AND REFERENCES,

PLATE CXXVIII.

Figures magnified 112 diameters.

	PAGE.
Figs. 1, 2, 4. VAUCHERIA GEMINATA, sections of true plant,	152
Figs. 3, 5. " " Var. <i>racemosa</i> ,	152
Fig. 6. VAUCHERIA, an abnormal but not rare form of fruiting,	152
" 7. A gall produced by a Rotifer; a Rotifer's nest,	152
" 8-10. VAUCHERIA HAMATA; California specimens,	152







EXPLANATIONS AND REFERENCES,

PLATE CXXIX.

Figures magnified 112 diameters.

	PAGE.
Fig. 1-5. VAUCHERIA TERRESTRIS, parts of fruiting filaments, variously developed with one, two or more oogonia on a peduncle, . . .	153
" 6-8. " " " oospores germinating, . . .	153
" 9-11. VAUCHERIA TUBEROSA, Var. <i>intermedia</i> , . . .	154
" 12-14 " " " <i>minor</i> , . . .	154









EXPLANATIONS AND REFERENCES,

PLATE CXXX.

Figures magnified 62 diameters.

	PAGE.
Fig. 1. VAUCHERIA TUBEROSA, type-form,	154
“ 1-5. Dichotomous branching and constricted basal ends of branchlets,	154
“ 6-10. Lateral stolons variously shaped ; the latter two tuberose,	154






EXPLANATIONS AND REFERENCES,

PLATE CXXXI.

Figures variously magnified.

		PAGE
Figs. 1, 2.	BOTRYDIUM GRANULATUM, twice the natural size,	155
Fig. 3.	“ “ two empty cells, the tops collapsed,	155
“ 4-7.	“ “ cells more highly magnified, showing the subterranean radicles,	155
“ 8.	“ “ cell-contents passing down into the radicles,	155
“ 9.	“ “ radicles developing zoospores,	155
“ 10.	“ “ cell-contents scattering zoömicrogonidia,	155
“ 11.	“ “ zoospores (sexual) highly magnified,	155
“ 12.	“ “ asexual macrospores,	155
“ 13.	“ “ zoospores developing in the radicles,	155
“ 14.	“ “ radicle cells under greater magnification.	156
Figs. 15, 16.	“ “ zoospores 4-6 days old,	156
“ 17, 18.	“ “ “ 7-8 “ “	156
Fig. 19.	“ “ zoospores germinating in radicles,	156
“ 20.	“ “ zoospores developing into vegetative plants,	156
Figs. 21, 22.	“ “ asexual zoospores enlarging and producing sexual zoospores,	156
“ 23, 24.	“ “ sexual zoospores copulating, producing isospores, (fig. 25),	156
Fig. 26.	“ “ isospores after a few days, becoming stellate,	156






EXPLANATIONS AND REFERENCES,

PLATE CXXXII.

Figures magnified 250 diameters.

	PAGE.
Fig. 1-3. SPIROGYRA TENUISSIMA,	207
Figs. 4, 5. SPIROGYRA SPREEIANA,	208
“ 6, 7. SPIROGYRA INFLATA,	207
Fig. 8-10. SPIROGYRA QUADRATA,	208
“ 11. SPIROGYRA WEBERI,	208
Figs. 12, 13. SPIROGYRA GREVILLEANA,	209




EXPLANATIONS AND REFERENCES,

PLATE CXXXIII.

Figures magnified 250 diameters.

	PAGE.
Figs. 1, 2. SPIROGYRA WEBERI, Var. <i>elongata</i> ,	209
“ 3, 4. SPIROGYRA CALOSPORA,	209
Fig. 5-7. SPIROGYRA HASSALLII,	210
Figs. 8, 9. SPIROGYRA INSIGNIS,	210
“ 10, 11. SPIROGYRA HANTZSCHII,	211





EXPLANATIONS AND REFERENCES,

PLATE CXXXIV.

Figures magnified 250 diameters.

	PAGE.
Figs. 1, 2. SPIROGYRA MIRABILE, 211
“ 3, 4. SPIROGYRA FLAVESCENS, 211
Fig. 5-7. SPIROGYRA GRACILIS, 211
“ 8-13. SPIROGYRA VARIANS, 212
“ 14-17. SPIROGYRA QUININA, 213






EXPLANATIONS AND REFERENCES,

PLATE CXXXV.

Figures magnified 250 diameters.

	PAGE.
Figs. 1, 2. SPIROGYRA INSIGNIS, Var. <i>elongata</i> ,	210
" 3, 4. SPIROGYRA ADNATA,	220
" 5, 6. SPIROGYRA DECIMINA,	216
" 7, 8. SPIROGYRA LUTETIANA,	214
" 9, 10. SPIROGYRA LONGATA,	214
" 11, 12. SPIROGYRA DUBIA,	220






EXPLANATIONS AND REFERENCES,

PLATE CXXXVI.

Figures magnified 250 diameters

	PAGE.
Fig. 1-3. SPIROGYRA FLUVIATILIS,	216
“ 4-7. SPIROGYRA RIVULARIS,	220
Figs. 8, 9. SPIROGYRA NEGLECTA,	216
“ 10, 11. SPIROGYRA ORTHOSPIRA,	218




EXPLANATIONS AND REFERENCES,

PLATE CXXXVII.

Figures magnified 250 diameters.

	PAGE.
Fig. 1-4. SPIROGYRA MAJUSCULA, 221
Figs. 5, 6. SPIROGYRA BELLIS, 217
“ 7, 8. SPIROGYRA NITIDA, 217




EXPLANATIONS AND REFERENCES,


PLATE CXXXVIII.

Figures magnified 250 diameters.

	PAGE.
Fig. 1-4. SPIROGYRA SETIFORMIS, 219
Figs. 5, 6 SPIROGYRA MAXIMA, Var. <i>inequalis</i> , 218
“ 7, 8. SPIROGYRA JUGALIS, 219




1



2




3



4

5



5




EXPLANATIONS AND REFERENCES,

PLATE CXXXIX.

Figures magnified 250 diameters.

		PAGE.
Figs. 1, 2.	SPIROGYRA BELLIS, compare Plate CXXXVII, figs. 5, 6,	. 217
“ 3, 4.	SPIROGYRA MAXIMA, see var. Plate CXXXVIII, figs. 5, 6,	218
“ 5, 6.	“ “ two spores germinating, 218
Fig. 7-9.	SPIROGYRA CONDENSATA, Var. <i>Rusbyi</i> ,	215




EXPLANATIONS AND REFERENCES,

PLATE CXL.

Figures magnified 200 diameters.

	PAGE.
Fig. 1-3. SPIROGYRA CRASSA, 219
“ 4-7. SPIROGYRA FUSCO-ATRA, 215
Figs. 8, 9. SPIROGYRA PARVISPORA, 221








EXPLANATIONS AND REFERENCES,

PLATE CXLI.

Figures magnified 250 diameters.

	PAGE.
Figs. 1, 2. SPIROGYRA SUBEQUA,	217
“ 3, 4. SPIROGYRA SUBSALSA,	212
Fig. 5-7. SPIROGYRA PUNCTATA,	215
“ 8-10. SIROGONIUM STICTICUM,	222





EXPLANATIONS AND REFERENCES,


PLATE CXLII.

Figures magnified 250 diameters.

	PAGE.
Figs. 1, 2. SPIROGYRA COMMUNIS, 213
“ 3, 4. SPIROGYRA JUERGENSEI,	213
“ 5, 6. SPIROGYRA MAJUSCULA, conjugated with <i>S. calospora</i> , 221
“ 7, 8. SPIROGYRA, rootlets, <i>S. quadrata</i> ,	208
“ 9, 10. SPIROGYRA, stellate nuclei, see figs. 8, 9, 210








EXPLANATIONS AND REFERENCES,
4
 PLATE CXLIII.

Figures magnified 300 diameters.

				PAGE.
Fig. 1-3.	ZYGNEMA LEIOSPERMUM,			222
“ 4-6.	ZYGNEMA INSIGNE,			223
Figs. 7, 8.	ZYGNEMA STELLIUM,			223
“ 9, 11.	“	forma <i>Vaucheria</i> ,		223
“ 12, 13.	“	“ <i>tenue</i> ,		223
“ 14, 15.	“	“ <i>subtile</i> ,		223
“ 16, 17.	“	“ <i>stagnale</i> ,		223





EXPLANATIONS AND REFERENCES,

PLATE CXLIV.

Figures magnified 300 diameters.

	PAGE.
Figs. 1, 2. ZYGNEMA CRUCIATUM,	224
Fig. 3-7. ZYGNEMA PURPUREUM,	224
“ 8. ZYGNEMA, spores germinating in cells,	224
“ 9. ZYGNEMA ANOMALUM,	224
“ 10-12. “ “ Var. <i>crassum</i> ,	224
“ 13. “ “ “ “ fertile,	225









EXPLANATIONS AND REFERENCES,

PLATE CXLV.

Figures magnified 300 diameters.

	PAGE.
Figs. 1, 2. ZYGOGONIUM PECTINATUM,	225
Fig. 3. ZYGOGONIUM AEQUALE,	226
Figs. 4, 5. ZYGOGONIUM DECUSSATUM,	226
“ 6, 7. ZYGOGONIUM AGARDHII,	227
Fig. 8. ZYGOGONIUM RALESII,	227
“ 9. ZYGOGONIUM PARVULUM,	227






EXPLANATIONS AND REFERENCES,

PLATE CXLVI.

Figures magnified 300 diameters.

		PAGE.
Figs. 1, 2.	MOUGEOTIA SPHAEROCARPA,	227
Fig. 3.	MOUGEOTIA MINNESOTENSIS,	228
“ 4.	MOUGEOTIA DIVARECATA,	228
“ 5.	MOUGEOTIA DELICATULA,	228
“ 6-9.	MOUGEOTIA GLYPTOSPERMA,	229




EXPLANATIONS AND REFERENCES.

PLATE CXLVII.

Figures magnified 300 diameters.

	PAGE.
Fig. 1. MESOCARPUS CRASSA,	230
Figs. 2, 3. MESOCARPUS SCALARIS,	230
Fig. 4. MESOCARPUS MACROSPORA,	230
“ 5. MESOCARPUS ROBUSTUS,	231
“ 6. MESOCARPUS RECURVUS,	231







EXPLANATIONS AND REFERENCES,

PLATE CXLVIII.

Figures magnified 300 diameters.

	PAGE.
Figs. 1, 2. MESOCARPUS NUMMULOIDES, 231
“ 3, 4. MESOCARPUS PARVULUS, 230
Fig. 5. MOUGEOTIA VERRUCOSUS, 229
“ 6. MESOCARPUS PARVULUS, Var.,
“ 7-10. MESOCARPUS RADICANS, 231
“ 11-15. PLAGIOSPERMUM TENUE, 233






EXPLANATIONS AND REFERENCES,


PLATE CXLIX.

Figures magnified 300 diameters.

	PAGE.
Fig. 1-5. GONATONEMA VENTRICOSUM,	233
“ 6-7. PLEUROCARPUS COLUMBIANUS,	232
Figs. 8, 9. PLEUROCARPUS MIRABILIS, ordinary sterile form,	232
Fig. 10-13. “ “ fruiting specimens,	232
“ 14-15. “ “ sterile filaments,	232








EXPLANATIONS AND REFERENCES,

PLATE CL.

Figures magnified 300 diameters.

	PAGE
Fig. 1-5. STAUROSPERMUM CAPUCINUM,	234
“ 6-8. STAUROSPERMUM QUADRATUM,	234
Figs. 9, 10. STAUROSPERMUM VIRIDE,	234
“ 11, 12. CRATERSPERMUM LETEVIRENS,	235




EXPLANATIONS AND REFERENCES,


PLATE CLI.

Figures magnified from 100-400 diameters.


		PAGE.
Fig. 1-4.	VOLVOX GLOBATOR, different stages of growth under low magnification; the latter with encysted resting spores,	158
“ 5.	Part of a coenobium of <i>Sphaerosira volvox</i> , Willn. Merely a condition of <i>Volvox globator</i> ,	159
“ 6.	Protoplasmic mass from the same under high magnification,	159
Figs. 7, 8.	The mass dividing into two, and four parts,	159
Fig. 9.	Dividing into thirty-two parts each with cilia, shown in lateral view,	159
“ 10.	A discoid family revolving in the mother sphere or apart from it, but in close proximity,	159
“ 11.	STEPHANOSPHAERA PLUVIALIS, a globose, vertical view,	163
“ 12.	Lateral view with fusiform primordial cells,	163
“ 13.	Formation of macrogonidia,	163
“ 14.	Division of macrogonidia,	163
“ 15.	GONIUM PECTORALE; lateral view of coenobium,	163
Figs. 16, 17.	Two coenobia in front view,	163
Fig. 18.	A coenobium before division,	163




1




2




3




4



5



6




7




8




9




10




11



12



13




14




15



16



17







EXPLANATIONS AND REFERENCES,

PLATE CLII.

	PAGE.
Fig. 1. VOLVOX GLOBATOR, a coenobium, showing the fine threads connecting the gonidia,	158
“ 2. Enlarged view of the connecting threads,	158
“ 3. An ideal section of the wall of a coenobium,	158
“ 4. Coenobium with three stellate resting spores,	158
“ 5. VOLVOX STELLATUS, Ehrb., a stellate oogonium or resting spore,	158
“ 6. An exaggerated ideal of a section of a coenobium,	158
“ 7. A complete antheridium under high magnification,	158
Figs. 8, 9. Swarming spermatozoids,	158
Fig. 11-20. EUDORINA STAGNALE,	160
“ 11. A coenobium of sixteen cells, normal vegetative condition,	160
“ 12. Cells dividing,	160
“ 13. A matured coenobium, nearly ready to scatter the young spheres,	160
“ 14-21. Spheres in different stages of development,	160
Figs. 22, 23. Spermatozoids,	160






EXPLANATIONS AND REFERENCES,

PLATE CLIII.

Figures mostly magnified 500 diameters.

	PAGE.
Fig. 1-5. <i>PANDORINA MORUM</i> , forms most frequent,	161
Figs. 6, 7. Coenobia with more numerous cells densely clustered,	161
“ 8, 9. Very fugacious forms, a variety or a distinct species.	161
Fig. 10. A matured coenobium with young spheres in condition for independent existence	161
“ 11-26. <i>CHLAMYDOCOCCUS PLUVIALIS</i>	164
Figs. 11, 12. Two cells of normal condition,	165
Fig. 13-26. Cells of polymorphous resting forms,	165
“ 27-29. Zoospores without envelope,	165
“ 30-32. Zoospores enveloped in a membrane,	165
“ 33. Commencement of a division of a green cell,	165
“ 34. Division completed.	165
“ 35-51. Numerous multiform conditions transpiring in the life of this little plant. For description of each see page,	166





EXPLANATIONS AND REFERENCES,

PLATE CLIV.

Figures magnified 500 diameters.

	PAGE.
Figs. 1, 2. CHLAMYDOCOCCUS NIVALIS, simple cells,	166
‘ 3, 4. Cells dividing into two and into four parts,	166
“ 5, 6. CHLAMYDOMONAS TINGENS, a variety of sizes of macrogonidia,	167
‘ 7, 8. CHLAMYDOMONAS PLUVIALE, the normal appearance,	167
“ 9, 10. CHLAMYDOMONAS HYALINA,	168
Fig. 11-14. HYDRODICTYON UTRICULATUM, parts of coenobia of various degrees of development,	170
Figs. 15, 16. First stages of development,	170
Fig. 17. Cell with microgonidia, in part escaping through a rupture of the membrane, highly magnified,	170
“ 18. Part of cell with macrogonidia, highly magnified,	170
“ 19. Free macrogonidia, more highly magnified.	170
“ 20. Zoospores or macrozoogonidia,	170







EXPLANATIONS AND REFERENCES,

PLATE CLV.

	PAGE.
Fig. 1-8. <i>EUGLENA VIRIDIS</i> , Ehrb., normal forms, capable of polymorphic contours,	161
“ 9-15. Cells contracting and coming to rest forming encysted spores,	162
“ 16-19. Advanced stages of encysted spores dividing into two, and into four parts,	162
“ 20. Encysted forms often occurring in countless numbers,	162
Figs. 21, 22. Sporangial bodies with escaping microzoogonidia,	162
“ 23, 24. <i>CHLAMYDOMONAS PULVIUSCULUS</i> , Ehrb.,	168
Fig. 25. An encysted form divided into two.	168
Figs. 26, 27. A similar form divided into four parts,	168
Fig. 28. Cell divided, forming new encysted cells,	168
“ 29. Two red resting cells,	168
“ 30. Macrozoospore; contents dividing,	168
“ 31. A cell with contents divided and developing microzoogonidia,	168
“ 32-35. <i>WOLFFIA COLUMBIANA</i> , Schleiden.	
A plant which has no claim to a place in this volume; it is admitted as an illustration. It is the smallest of phenogamous plants, and is a form very apt to deceive in the search for fresh-water algae. It is found in wet, marshy, places, not frequent, but in quantity when it does occur. Fronds are rootless, floating as small grains, about the size of poppy seed, in shallow, stagnant waters. Cells broadly oval. Multiplication takes place by the formation of a daughter cell in the maternal frond, which opens at one end and the young slides out. Figs. 32-35, represent progressive stages of proliferation.	
“ 36-40. <i>MICHOCOCCUS CONFERVICOLA</i> , matured plants and young developing forms,	200
“ 41. <i>COSMOCLADIUM PULCHELLUM</i> , figure transcribed from Rabenhorst; hitherto not found in the United States.	






EXPLANATIONS AND REFERENCES,

PLATE CLVI.

Figures magnified 500 diameters.

	PAGE.
Fig. 1-3. COELASTRUM MICROSPORUM,	170
“ 4. Variety <i>speciosum</i> ,	170
“ 5. COELASTRUM CAMBRICUM,	170
“ 6-10. SORASTRUM SPINULOSUM, coenobia of various forms, differing according to age,	171
Figs. 11, 12. SCENEDESMUS CAUDATUS, typical form.	172
“ 13, 14. Smaller forms,	172
Fig. 15. Var. <i>horridus</i> ,	172
Figs. 16, 17. SCENEDESMUS ANTENNATUS, Var. <i>rectus</i> ,	172
Fig. 18-21. SCENEDESMUS DIMORPHUS, different forms,	173
“ 22-24. SCENEDESMUS OBTUSUS, various forms,	173
“ 25-26. SCENEDESMUS ACUTUS, two fronds,	173
“ 27. SCENEDESMUS ROTUNDATUS,	174
“ 28. DICTYOSPHAERIUM RENIFORME,	186
Figs. 29, 30, DICTYOSPHAERIUM EHRENBERGIANUM,	186
Fig. 31. A family after Naegeli's figure,	186




EXPLANATIONS AND REFERENCES,

PLATE CLVII.

Figures magnified 500 diameters.

	PAGE.
Figs. 1, 6. <i>SCIADIUM ARBUSCULA</i> ,	174
“ 1, 2. Young cells,	174
Fig. 3. Commencement of the first generation of daughter cells,	174
“ 4. Farther progress of growth,	174
Figs. 5, 6. A second and a third generation,	174
“ 7, 8. <i>SCIADIUM GRACILIPES</i> from a specimen from a pool, Minnesota,	175
Fig. 9-11. <i>STAUROGENIA CRUCIATUM</i> , three different forms,	171
“ 12. <i>DICTYOSPHAERIUM HITCHCOCKII</i> , frequent in ponds, New Jersey,	186







EXPLANATIONS AND REFERENCES,

PLATE CLVIII.

Figures magnified 500 diameters.

	PAGE.
Figs. 1, 2	OPHIOCYTIUM CUSPIDATUM, after specimens from Massachusetts
	and from California, 176
Fig. 3-7.	OPHIOCYTIUM CAPITATUM, various forms, 176
“ 8-14.	OPHIOCYTIUM COCHLEARE, usual appearances, 175
“ 15-18.	OPHIOCYTIUM CIRCINATUM, from Minnesota, 176






EXPLANATIONS AND REFERENCES,

PLATE CLIX.

Figures magnified 500 diameters.

		PAGE.
Fig. 1.	CHARACIUM SESSILE five cells, 177
" 2.	CHARACIUM ACUTUM, shortly stipitate, 177
" 3.	CHARACIUM PRINGSHEIMII, larger and longer stipitate, 177
" 4.	CHARACIUM NÆGELII, 178
" 5.	CHARACIUM AMBIGUUM, 177
" 6.	CHARACIUM HETEROMORPHUM, 178
" 7-10.	POLYEDRIUM TRIGONUM, Var. <i>tetragonum</i> , 184
" 11-14.	POLYEDRIUM GIGAS, type-form, 184
" 15-18.	POLYEDRIUM TRIGONUM, Var. <i>bifurcatum</i> , 184
" 19-23.	POLYEDRIUM ENORME, 184
" 24.	POLYEDRIUM MUTICUM, 185
" 25-27.	POLYEDRIUM TRIGONUM, Var. <i>minus</i> , 184
" 28-34.	POLYEDRIUM MINIMUM, some in vegetative condition, others discharging gonidia, 185
" 35-38.	POLYEDRIUM LONGISPINUM, 185







EXPLANATIONS AND REFERENCES,

PLATE CLX.

Figures magnified 500 diameters.

	PAGE.
Figs. 1, 2. RHAPHIDIUM CONVOLUTUM, (<i>Selenastrum</i>),	198
“ 4, 5. “ “ small form,	199
“ 6, 7. RHAPHIDIUM POLYMORPHUM, Var. <i>fulcatum</i> ,	187
“ 8, 10, “ “ in different stages of growth,	198
“ 11, 12. “ “ Var. <i>contortum</i> ,	198
Fig. 13-15. “ “ “ “ from Florida,	198
“ 16-21. “ “ fruiting forms,	198
Figs. 22, 23. “ “ Var. <i>aciculare</i> ,	197
“ 24, 25 “ “ Var. <i>sigmoideum</i> ,	198
“ 26, 27. RHAPHIDIUM BRAUNII,	198
“ 28, 29. RHAPHIDIUM CONVOLUTUM, Var.,	199
Fig. 30-35. DIMORPHOCOCCUS CORDATUS, coenobia in different stages of growth,	199
“ 36-38. “ “ simple groups of cell in early stage of development,	199
















EXPLANATIONS AND REFERENCES,

PLATE CLXII.

Figures magnified 500 diameters.

		PAGE.
Fig. 1.	PROTOCOCCUS VESTITUS, small, loose family,	183
" 2.	PROTOCOCCUS VIRIDIS, Var., the more unusual condition,	181
" 3.	" " Var. <i>angulosa</i> ,	181
" 4.	" " Var. <i>dissectus</i> ,	181
" 5.	" " Var. <i>miniatus</i>	181
" 6.	PALMELLA HYALINA,	193
" 7.	SCHIZOCHLAMYS GELATINOSA, A. Br. (a condition of <i>Tetraspora</i>),	192
" 8.	PALMELLA BOTRYOIDES,	194
" 9.	SCHIZOCHLAMYS DECORTICANS,	192
" 10.	PALMELLA MINIATA,	193
" 11.	PALMELLA MOOREANA,	193
" 12.	PALMELLA MUCOSA,	192
" 13.	PALMELLA UVAEFORMIS,	193
" 14.	PALMELLA MUCOSA, Var.,	192







EXPLANATIONS AND REFERENCES,

PLATE CLXIII.

Figures magnified 500 diameters.

	PAGE.
Fig. 1-5.	PROTOCOCCUS VIRIDIS, Vars. <i>glomeratus</i> , <i>dissectus</i> and <i>humicola</i> , showing early stages of development of filaments of <i>Ulothrix</i> , . . . (138), 181
" 6.	PROTOCOCCUS INFUSIONUM, 182
" 7.	A form frequent in pools,
Figs. 8, 9.	ULOTHRIX PARIETINA, developing from <i>Protococcus</i> , . . . 138
Figs 10, 11.	Early stages of development of <i>Stigeoclonium</i> from <i>Protococcus</i> , compare Plate XCVI,
" 12, 13.	NEPHROCYTIUM AGARDHIANUM, 197
" 14, 15, 17.	" " younger forms, 197
" 16, 18, 19.	NEPHROCYTIUM NAEGELII, 197








EXPLANATIONS AND REFERENCES,

PLATE CLXIV.

Figures variously enlarged.

			PAGE.
Figs. 1, 2.	HYDRURUS FETIDUS, Var. <i>penicillatus</i> ,	about half the natural size,	187
“ 3, 4.	“ “	small section of thallus much enlarged, showing the scattered cells,	189
“ 5, 6.	“ “	two young plants.	189
Fig. 7-9.	“ “	three cells taken from matured branchlets (fig. 4), greatly magnified,	189
“ 10-12.	“ “	progressive conditions of development of cells (gonidia), after Rostafinski,	189
“ 13.	“ “	Cells dividing,	189
Figs. 14, 15.	HYDRURUS IRREGULARIS, forma <i>confusa</i> ,		189
“ 16, 17.	HYDRURUS DUCLUZELII,		189






EXPLANATIONS AND REFERENCES,

PLATE CLXV.

Figures under different magnification.

	PAGE.
Fig. 1-4. TETRASPORA BULLOSA, thallus about natural size; the cells magnified,	190
Figs. 5, 6. " " thallus less than natural size,	190
" 7, 8. TETRASPORA CYLINDRICA, thallus one-half the natural size; cells magnified,	190
Fig. 9. TETRASPORA LUBRICA, about half the natural size,	191
Figs. 10, 11. " " Var. <i>lacunosa</i> , half natural size; cells magnified,	191
Fig. 12. TETRASPORA EXPLANATA,	191














EXPLANATIONS AND REFERENCES,

PLATE CLXVI.

Figures variously magnified.

	PAGE.
Fig. 1. PORPHYRIDIVM CRVENTVM, a very small portion of a thallus,	194
“ 2-8. GLOEOCYSTIS AMPLA, variety of cells,	196
“ 9-15. GLOEOCYSTIS VESICULOSA,	196
“ 16-18. GLOEOCYSTIS RUFESCENS,	196
“ 19-21. GLOEOCYSTIS RUPESTRIS,	196



10

14

17

18

18

18


20

21

EXPLANATIONS AND REFERENCES,

PLATE CLXVII.

			PAGE.
Fig. 1.	EREMOSPHAÆREA VIRIDIS,	a cell in normal condition shedding	
		its membranous envelope,	200
' 2.	"	" a cell dividing,	200
Figs. 3, 4.	"	" a cell dividing into eight sister cells,	200
Fig. 5-11.	"	" different conditions of cell multiplication forming of gonidia, resting spores, etc.,	200










EXPLANATIONS AND REFERENCES,

PLATE CLXVIII.

		PAGE.
Figs. 1, 2	CALOTHRIX ORSINIANA, two filaments,	236
“ 3, 4.	CALOTHRIX DILLWYNI,	237
Fig. 5.	CALOTHRIX GYPSOPHYLA,	237
Figs. 6, 7.	“ “ smaller form,	237








EXPLANATIONS AND REFERENCES,

PLATE CLXX.

	PAGE.
Figs. 1, 2. CALOTHRIX RADIOSA, two plants drawn on a small scale,	238
“ 3, 4. CALOTHRIX HOSFORDII, a variety which appears to hold a position between <i>C. radiosa</i> and <i>Calothrix Meneghiniana</i> ,	238
Fig. 5-7. CALOTHRIX MENEGHINIANA, three variously developed forms,	238
Figs 8, 9. Young conditions of growth of the plants of this plate,	238








EXPLANATIONS AND REFERENCES,

PLATE CLXXI.

	PAGE.
Fig. 1-4. MASTIGONEMA AERUGINEUM, a distinct mode of fruiting, the basal cells developing macrogonidia; from Florida,	. 242
Figs. 5, 6. Filaments from a floating mass; basal heterocysts torn off; brownish color,	. 242
Fig. 7. Ordinary condition of Florida specimens,	. 242
Figs. 8, 9. Three branching forms from a gelatinous coating on the sides of a pump trough,	. 241









EXPLANATIONS AND REFERENCES,

PLATE CLXXII.

		PAGE
Fig. 1.	CALOTHRIX LACUCOLA,	239
“ 2.	CALOTHRIX CRUSTACEUM, a single plant separated from an in- crustation, under higher magnification,	240
“ 3.	“ “ a section of a shining blackish frond,	240
Figs. 4, 5.	“ “ the gray crustiferous form, natural size,	240
Fig. 6.	“ “ section of a more advanced growth,	240
“ 7.	“ “ matured, older plant; ends of tri- chomes fibrillose,	240





EXPLANATIONS AND REFERENCES,

PLATE CLXXIII.


	PAGE.
Figs. 1, 2, 3. CALOTHRIX GRACILIS, differs from <i>C. Dillwynii</i> mainly in size,	237
“ 4, 5. CALOTHRIX AERUGINEUM, young and maturer forms from Florida,	242
Fig. 6-8. MASTIGONEMA SEJUNCTUM, younger isolated trichome; section of matured plant, and full grown plant,	243
“ 9. “ “ a small tuft,	243




EXPLANATIONS AND REFERENCES,

PLATE CLXXIV.

		PAGE.
Fig. 1.	MASTIGONEMA HALOS, the plant enlarged 250 diameters,	. 242
" 2.	" " under low magnification, . . .	243
" 3-9.	MASTIGONEMA AERUGINEUM, in varieties, 241
" 3-4.	" " distinct forms with twin hetero- cysts, 241
" 5-7.	" " with pyriform heterocysts, 241
" 8.	" " spherical, single heterocysts (<i>M.</i> <i>fibrosa</i>), 244, 241
" 9.	MASTIGONEMA ELONGATUM, 243






EXPLANATIONS AND REFERENCES,

PLATE CLXXV.

			PAGE.
Fig. 1-3.	MASTIGONEMA AERUGINEUM.	developing from spores; different stages of advancement	241
“ 4.	“	“ developing <i>tolypothrix</i> ,	241
Figs. 5, 6.	“	“ other developing forms,	241
Fig. 7.	“	“ developing conditions; short flagelliform, then elongating changing to linear,	241
“ 8.	“	“ positions of trichomes changed from younger cluster, fig. 9,	241
“ 9.	“	“ younger condition than fig. 8,	241
“ 10.	“	“ condition surrounded by many spores, after development of filaments,	241
“ 11.	“	“ an occasional condition of a filament under higher power,	241
Fig. 12-14.	“	“ a larger form from New Jersey, others, fig. 14, from Florida,	241
“ 15-17.	“	“ more scattered trichomes,	241








EXPLANATIONS AND REFERENCES,

PLATE CLXXVI.

	PAGE.
Fig. 1. MASTIGONEMA AERUGINEUM; a form parasitic on <i>Lymnobia</i> <i>Wollei</i> , and probably a young development of it,	242
“ 2-4, “ “ another form indicating the same mode of branching as Plate CLXXI, fig. 8-10,	241
“ 5. ISACTIS CAESPITOSA, small section of a thallus,	245
Figs. 6, 7. MASTIGONEMA VELUTINA, young condition of <i>Lymnobia aestuarii</i> .	242
Fig. 8. “ “ early development of the last,	242
“ 9-11. “ “ differing principally in having inter- stitial heterocysts,	242








EXPLANATIONS AND REFERENCES,

PLATE CLXXVII.

Figures mostly magnified 250 diameters.

		PAGE.
Figs. 1, 2.	MASTIGONEMA forms, somewhat distinct in the heterocysts and in the fibrillose sheaths,	242
“ 3, 4.	“ more common forms,	242
“ 5, 6.	ISACTIS FLUVIATILIS, two small sections of crust on rocks, .	245
Fig. 7-10.	“ “ isolated trichomes separated from the crust, 245	
“ 11.	MASTIGONEMA AERUGINEUM, elongated form,	242
“ 12.	“ “ turgid specimens,	242
“ 13.	“ “ heterocysts wider than the tri- chomes,	242
“ 14.	CALOTHRIX CRUSTACEUM, section of thallus representing the <i>Zonotrichia</i> condition,	240
“ 15.	“ “ natural size of thalli, often many clustered together; fig. 14 is a section of one of these magnified,	240






EXPLANATIONS AND REFERENCES,

PLATE CLXXVIII.

Figures magnified 250 diameters

	PAGE.
Fig. 1-3. <i>ISACTIS CAESPITOSA</i> , from specimens, Vermont and New Jersey,	245
“ 4-7. <i>GLOEOTRICHIA NATANS</i> , progressive stages of the growth of the sheath of a trichome,	246
“ 8-12. Progressive stages of another form,	246
Figs. 13, 14. Trichomes fertile, and fading, in the same thallus,	246
Fig. 15-19. Smaller, young, and older thalli, natural size,	246
“ 20. A form from Colorado,	246






EXPLANATIONS AND REFERENCES,

PLATE CLXXIX.

		PAGE.
Fig. 1.	RIVULARIA DURA, transverse section of a large thallus,	. 249
“ 2.	“ “ section of a semi-thallus, developing spores,	249
“ 3.	“ “ section of a semi-thallus, with trichomes withering,	249
“ 4.	RIVULARIA ECHINULATA, one of a floating mass of thalli,	. 249
“ 5.	“ “ natural size of thalli,	249
“ 6.	GLOEOTRICHIA PISUM, thalli of natural size, 247
“ 7.	“ “ peculiar condition; endochrome passing down into the basal heterocysts, 247
Figs. 8, 9.	“ “ small sections of other thalli not empty- ing in the same manner, 247
Fig. 10.	GLOEOTRICHIA NATANS, two trichomes, apparently in a state of dissolution, 247
“ 11.	“ “ trichomes with long spores, narrow sheath; gonidia passing out at rup- tured ends,	247




EXPLANATIONS AND REFERENCES,

PLATE CLXXX.

Figures magnified 250 diameters.

	PAGE.
Fig. 1-3. TOLYPOTHRIX DISTORTA, usual form,	263
“ 4. “ “ forma <i>magma</i> ,	263
Figs. 5, 7. TOLYPOTHRIX PULCHRA,	264
Fig. 8-10. TOLYPOTHRIX RAVENELII, from Florida specimens,	265
“ 11-13. TOLYPOTHRIX RUPESTRIS, Pennsylvania,	265
“ 14-16. TOLYPOTHRIX FLACCIDA, parts of three filaments,	265







EXPLANATIONS AND REFERENCES,

PLATE CLXXXI.

Figures magnified 250 diameters.

	PAGE.
Fig. 1-4. TOLYPOTHRIX AEGAGROPILA or MUSCICOLA,	263
" 5-7. TOLYPOTHRIX TENUIS, ponds, Pennsylvania,	265
" 8-11. TOLYPOTHRIX TRUNCICOLA, frequent on trunks of trees, South Carolina,	266
" 12-15. PLECTONEMA MIRABILE, typical forms, limestone springs, Penn- sylvania,	266








EXPLANATIONS AND REFERENCES,

PLATE CLXXXII.

Figures magnified 250 diameters.

	PAGE.
Fig. 1-3. SCYTONEMA MYOCHROUS; three forms,	252
“ 4. SCYTONEMA TOLYPOTRICHOIDES, a normal vegetative condition, 250	
“ 5. “ an older plant,	251
Figs. 6, 7. “ part of filament with secondary branches,	251
Fig. 8. “ a young form just developed from a <i>Nostoc</i> , (255), 251	
Figs. 9, 10. “ <i>Nostoc</i> forms developing <i>Scytonema</i> , (255), 251	
Fig. 11-13. “ less advanced stage of development,	251




EXPLANATIONS AND REFERENCES,

PLATE CLXXXIII.

Figures magnified 250 diameters.

	PAGE.
Figs. 1, 2. SCYTONEMA CATARACTA, from Rocky Mountain specimens,	. 252
“ 3, 4. “ “ from Niagara specimens,	. 252
Fig. 5-7. SCYTONEMA CALOTRICHOIDES, three specimens from submerged sticks, pond, Brown's Mill, New Jersey, 251
“ 8-10. SCYTONEMA NATANS, frequent in ponds, New Jersey,	. 251
“ 11-13. SCYTONEMA NAEGELII, typical forms, 252




EXPLANATIONS AND REFERENCES,

PLATE CLXXXIV.

Figures magnified 250 diameters.

			PAGE.
Figs. 1. 3.	SCYTONEMA	NOTARISII, normally developed plants,	254
“ 4, 5.	“	“ younger growths,	255
Fig. 6-9.	“	“ more juvenile forms,	255
Figs. 10, 11.	“	“ gonidia developing,	255
“ 12, 13.	“	“ large cysts containing gonidia,	255
Fig. 14.	“	“ young forms,	255
Fig. 16-19.	SCYTONEMA	CASTELLII, parts of matured plants,	255
“ 20-21.	“	“ parts of two older filaments, the articulations, short-discs, sliding out of the sheaths, scattering microgonidia, which in turn increase in size from (21) — (23) — (24), then redivide ejecting macrogonidia and producing a new cyst, <i>Microcystis</i> , <i>Gloeothecae</i> , and other forms,	255
“ 24-29.	“	“ cysts developing young plants,	255
“ 30-33.	“	“ young plants,	255
“ 34-40.	“	“ <i>Nostoc</i> and <i>Scytonema</i> forms,	255




EXPLANATIONS AND REFERENCES,

PLATE CLXXXV.

Figures magnified 250 diameters.

	PAGE.
Fig. 1-4. SCYTONEMA GRACILE, after specimens from rocky shores of Morris Pond, New Jersey,	253
“ 5-7. SCYTONEMA TURFOSUM, medium size filaments from exposed soil,	253
“ 8-10. SCYTONEMA CINCINNATUM, ponds, Pennsylvania and other States,	254





EXPLANATIONS AND REFERENCES,


PLATE CLXXXVI.

Figures magnified 250 diameters.

			PAGE.
Fig. 1-4.	SCYTONEMA CORTEX, forma <i>Ravenelii</i> ,	.	257
Figs. 5, 6.	“ “	young conditions from bark of trees,	257
Fig. 7-9.	“ “	forma <i>saxicola</i> , a rock form,	257
“ 10-14.	SCYTONEMA INTERTEXTUM, five distinctive parts of filaments,		258
Figs. 15, 16.	“ “	microgonidia, in developing conditions,	258
“ 17, 18.	“ “	young conditions,	258









EXPLANATIONS AND REFERENCES,

PLATE CLXXXVII.

Figures magnified 250 diameters.

		PAGE.
Fig. 1-3.	SCYTONEMA MIRABILIS, parts of three filaments,	255
“ 4.	“ “ microgonidia developed from the cells of the filaments,	256
“ 5,	“ “ the microgonidia enlarging, forming cysts— <i>Microcystis</i> forms,	.
“ 6.	“ “ the <i>Microcystis</i> forms developing <i>Gloeocapsa</i> forms,	256
“ 7-9.	“ “ <i>Nostoc</i> forms from the last,	.
“ 10-15.	“ “ <i>Nostoc</i> filaments having developed sheaths,	256
Figs. 16, 17.	“ “ imperfectly developed or faded filaments,	256
Fig. 18-21.	SCYTONEMA CORTEX, four samples of larger branching forms,	256
Figs. 22, 23.	“ “ two broken off ends, with cells singly, or in series sliding out,	257
Fig. 24.	“ “ two cells enlarging, developing macrogonidia,	257
Figs. 25, 26.	“ “ these in turn increase in size, producing first, <i>Microcystis</i> forms,	257
Fig. 27.	“ “ then <i>Nostoc</i> forms,	257
Figs. 28, 29.	“ “ from these cells are developed young plants,	257
Fig. 30-33.	“ “ young <i>Nostoc</i> forms producing <i>Scytonema</i> ,	257, 255
“ 34-37.	“ “ young forms of <i>Scytonema</i> ,	257










EXPLANATIONS AND REFERENCES,

PLATE CLXXXIX.

Figures magnified 250 diameters.

		PAGE.
Fig. 1.	SYMPHYOSIPHON CASTELLI, a cluster of filaments, . . .	262
“ 2.	SYMPHYOSIPHON AMBIGUUM, often intermixed with other algae on moist rocks.	262
“ 3.	SYMPHYOSIPHON HOFMANNI, a simple cluster; not always so spreading,	262
“ 4.	SYMPHYOSIPHON BORNETIANUM, from specimens coating old bricks,	261
“ 5.	SYMPHYOSIPHON AUSTINII, a few filaments separated from an almost black stratum,	261
“ 6.	SYMPHYOSIPHON BORETIANUM, from clay cliffs, South Carolina,	261
“ 7.	SYMPHYOSIPHON HIRTULUS, the normal growth on damp rocks, and moist compact soil,	261
“ 8.	“ “ filaments separated from clusters,	262






EXPLANATIONS AND REFERENCES,

PLATE CXC.

Figures magnified 250 diameters.

			PAGE.
Fig. 1-3.	SIROSIPHON PULVINATUS, macrogonidia, (<i>Gloeocapsa</i> cells),		269
“ 4-8.	“	“ early development from macrogonidia,	269
Figs. 9, 10.	“	“ two elongated filaments with a single series of cells,	269
“ 11, 12.	“	“ older forms with 2-6-8 series of cells.	269





EXPLANATIONS AND REFERENCES.

PLATE CXCI.

Figures magnified 250 diameters.

	PAGE.
Figs. 1, 2. SIROSIPHON PULVINATUS, forma <i>Alpinus</i> , two ends of filaments with young spine-like growths,	269
Fig. 3. A broken-off section of a filament, the cells bearing the micro- gonidia enlarging and sliding out of the sheath,	269
Figs. 4, 5. The same increasing in size,	.
“ 6, 7. Developing encysted cells, <i>Gloeocapsa</i> ; tegument breaks; cells are scattered,	269
Fig. 8-10. Cells pass through another cycle of development,	269
“ 11-14. Young plants developing from the cells (8-10),	269
Figs. 15, 16. Another form of microgonidia,	270
“ 17, 18. Advanced stages of development,	270
“ 19, 20. Gradually attain to the condition of <i>Gloeocapsa</i> ,	270
Fig. 21. SIROSIPHON CORALLOIDES, a single, rather large stem from a tomentose mat, growing on stones of shore of Green Pond, New Jersey,	270






EXPLANATIONS AND REFERENCES,

PLATE CXCII.

Figures magnified 250 diameters.

		PAGE.
Figs. 1, 2.	SIROSIPHON COMPACTUS, several characteristic filaments,	. 271
Fig. 3-5.	SIROSIPHON SCYTONEMATOIDES,	271
" 6.	" " " macrogonidia, 271
" 7-8.	HAPALOSIPHON?	
" 9-12.	SIROSIPHON CORALLOIDES, Var. <i>inflatus</i> , specimens from Mt. Mansfield, Vermont.	270





EXPLANATIONS AND REFERENCES,

PLATE CXCVIII.

Figures magnified 250 diameters.

	PAGE.
Figs. 1, 2. SIROSIPHON LIGNICOLA, two well developed forms,	273
“ 3, 4. “ “ an occasional appearance of a distorted growth,	273
Fig. 5-11. “ “ conditions of young developments,	273
Figs. 12, 13 SIROSIPHON ARGILLACEUS, two simple forms from a cluster,	273






EXPLANATIONS AND REFERENCES,

PLATE CXCIV.

Figures magnified 250 diameters.

	PAGE.
Fig. 1-3. SIROSIPHON OCELLATUS, three sections of branching filaments,	272
“ 4. Part of a broken filament,	272
Figs. 5, 6. Cells passing out and developing into larger forms,	272
Fig. 7-10. Young plants developing from the last,	272
“ 11-16. SIROSIPHON CRAMERI. from specimens from New Jersey answering the descriptions, but suspect, it may be merely a form <i>S. ocellatus</i> ,	272
“ 17-20. SIROSIPHON BRANDEGEEI, older and younger forms,	274
“ 21. Sporangial cells,	274
Figs. 22, 23. <i>Gloeocapsa</i> cells developed from the last condition,	274
Fig. 24-27. Conditions of young developing plants,	274




EXPLANATIONS AND REFERENCES,

PLATE CXCIV.

Figures magnified 250 diameters.

	PAGE.
Fig. 1-4. SCYTONEMA HEPPH,	260
“ 5. “ “ a broken-off filament,	260
Figs. 6, 7. “ “ the macrospores escaping and developing ; reproducing young plants,	260
“ 8, 9. “ “ young conditions,	260
Fig. 10. HAPALOSIPHON BYSSOIDEUS, in early stage,	276
Figs. 11, 12. “ “ more advanced,	276
“ 13, 14. HAPALOSIPHON TORULOSUS part of a faded filament,	276
Fig. 15. SIROSIPHON PULVINATUS, from West Virginia, very tuberclose, near Agardh's <i>S. mamillosum</i> ,	270
“ 16. “ “ specimen from dripping rocks West Virginia,	270




EXPLANATIONS AND REFERENCES,

PLATE CXCVI.

Figures magnified 250 diameters.


	PAGE.
Fig. 1. HAPALOSIPHON BREBISSENI, from specimens, Dennisville, N.J.,	276
“ 2-4. HAPALOSIPHON BRAUNII, parts of three filaments with branches shortened,	275
“ 5. “ “ small section, more highly magnified to show the process of transverse divi- sion of cells. As they enlarge they glide out of the filament, and then continue growth,	276
“ 6 “ “ producing encysted cells,	276
Figs. 7, 8. From these last <i>Chroococcus</i> cells are developed, also the follow- ing four,	276
“ 9, 10. <i>Palmagloea</i> cells, (<i>Mesotaenium</i>), see Desmids U. S.,	276
Fig. 11-15. <i>Nostochopsis</i> (Wood) forms,	276
Figs. 16, 17. <i>Nostocs</i> ,	276
Fig. 18. <i>Hormosiphon</i> ,	276
“ 19. A specimen developing secondary branchlets,	276
Figs. 20, 21. HAPALOSIPHON TENUISSIMUS, not infrequent, ponds, New Jersey, Pennsylvania, Minnesota, Florida,	277
“ 22, 23. HAPALOSIPHON FUSCESCENS, pond waters, New Jersey,	277



EXPLANATIONS AND REFERENCES,

PLATE CXCVII.


	PAGE.
Fig. 1. NOSTOC VERRUCOSUM, a frond,	285
“ 2. “ “ moniliform filaments from a frond,	285
Figs. 3, 4. NOSTOC COERULEUM, <i>N. minutissimum</i> , Kg.,	(279), 284
Fig. 5. NOSTOC PRUNIFORME, well developed forms,	284
Figs. 6, 7. “ “ filaments from a frond	284
Fig. 8. NOSTOC COMMUNE, a not infrequent form of thallus,	283
“ 9. Normal form of filaments,	283
Figs. 10, 11. Cells of filaments separated and developing young plants.	283
Fig. 12-15. Forms of newly developed plants,	283
Figs. 16, 17. Two young filaments, in part developed, in part unchanged, moniliform,	283
Fig. 18-20. NOSTOC SPHAERICUM, natural sizes as they vary with age,	283
“ 21-23. NOSTOC RUPESTRE, not separable from the last, but more fully de- veloped. Filaments primarily without sheaths,	283
Figs. 24, 25. A more matured frond, filaments have formed sheaths,	283
Fig. 26-30. Forms which constitute the <i>N. rupestre</i> , Kg., embracing also <i>N.</i> <i>macrosporum</i> , Menegh.; <i>N. coreaceum</i> , Vauch.; <i>N. verrucosum</i> , Vauch.; <i>N. tenuissimum</i> , Rab.; <i>N. microscopicum</i> , Carm.; <i>N. gregarium</i> , Hantzsch.; <i>Hormosiphon furfuraceus</i> , Kg., and others, all of which are merely different stages of growth,	283
“ 31. A multiplication of <i>Nostocs</i> ,	278
“ 32. NOSTOC FLAGELLIFORME, from specimens from Texas,	285
Figs. 33, 34. NOSTOC ALPINUM, Kg., <i>N. cristatum</i> , Bail.; <i>N. parmelioides</i> , Kg.,	285
Fig. 35. NOSTOC MUSCORUM,	282



EXPLANATIONS AND REFERENCES,

PLATE CXCVIII.

	PAGE.
Figs. 1, 2. ANABÆNA CUPRESSAPHILA, from trunks of cedar trees near water's edge, swamps, New Jersey,	289
" 3, 4. NODULARIA PALUDOSA, from specimens, Colorado and Pennsylv- vania,	291
" 5, 6. APHANIZOMENON FLOS-AQUAE, short sections with spores, .	291
" 7, 8. " " agglutinated filaments, .	291
" 9, 10. ANABÆNA, filaments, one in normal vegetative, moniliform con- dition and the other in process of cell division, 286, (298)	
Fig. 11-13. ANABÆNA, three progressive conditions of development, 286, (298)	
" 14-19. ANABÆNA, advanced stages of growth,	(286, 298)
" 20-23. ANABÆNA, FLOS-AQUAE, filaments of type-form,	287
" 24-26. " " Var. <i>circinalis</i> ,	287
Figs. 27, 28. ANABÆNA GIGANTEA, two short sections,	287
Fig. 29-32. ANABÆNA VARIABILIS,	287
" 33-35. SPHAEROZYGA HASSALLII, three filaments, specimens of many variable forms,	289
" 36. OSCILLARIA, filament resulting from the last,	289
Figs. 37, 38. SPHAEROZYGA POLYSPERMA, from ponds, pools, etc.,	289
" 39, 40. ANABÆNA OSCILLARIOIDES, filaments in different stages of de- velopment,	288




EXPLANATIONS AND REFERENCES,

PLATE CXCIX.

Figures magnified 250 diameters.


		PAGE.
Fig. 1.	SPHAEROZYGA SACCATA, a cluster of saccate thalli,	290
“ 2.	“ “ older detached floating forms,	290
“ 3-5.	“ “ specimens of filaments contained in the saccate thalli,	290
“ 6-8.	CYLINDROSPERMUM MACROSPERMUM, normal appearance of the filaments,	292
“ 9-10.	CYLINDROSPERMUM LIMNICOLA, after specimens from a ditch almost dried up. Some of the filaments appear to be veritable <i>Oscillaria</i> ,	292
“ 11.	“ “ cells of filaments indistinct, not constricted at joints; <i>Oscillaria</i> -like,	292
“ 12.	“ Wood's figure of <i>C. minutum</i> ,	292
“ 13.	“ Wood's figure of <i>C. flexuosum</i> ,	293
“ 14.	“ Wood's figure of <i>C. macrospermum</i> ,	292
“ 15.	ANABÆNA, FLOS-AQUÆ,	286
“ 16.	CYLINDROSPERMUM COMATUM, Wood's figure,	293
“ 17.	SPHAEROZYGA SMITHII; <i>Hilsia</i> , Kirch.,	290
Figs. 18, 19.	“ “ typical filaments,	290
Fig. 20-22.	“ “ filaments with two or three trichomes in a sheath,	290
Figs. 23, 24.	“ “ more advanced conditions of older filaments,	290




EXPLANATIONS AND REFERENCES,

PLATE CC.

	PAGE.
Figs. 1, 2. CHAMAESIPHON CONFERVICOLA,	295
Fig. 3. CHAMAESIPHON INCRUSTANS, young forms often densely placed, incrusting the plants upon which it is parasitic,	295
Figs. 4, 5. LYNGBYA CINCINNATA, from specimens from Mt. Mansfield, Vermont,	296
Fig. 6-8. LYNGBYA WOLLEI, short sections of filaments, often very abundant,	297
Figs. 9, 10. LYNGBYA MAJUSCULA, from brackish water, New York,	297
Fig. 11-16. LYNGBYA AESTUARII, specimens collected in numerous localities along the coast of New Jersey, and in Florida,	296








EXPLANATIONS AND REFERENCES,

PLATE CCI.

Figures magnified 250 diameters.

	PAGE.
Figs. 1, 2. <i>LYNGBYA NAVEANUM</i> , Pennsylvania specimens,	298
“ 3, 4. “ “ Florida specimens.	298
Fig. 5-10. <i>LYNGBYA OBSCURA</i> . Kg., specimens from Canada, Vermont, and Pennsylvania,	298
“ 11-13. <i>LYNGBYA MEMBRANACEA</i> , specimens from Charlotte, Vermont,	300
“ 14-20. <i>LYNGBYA VULGARIS</i> , frequent from Maine to Florida,	300
“ 21. <i>LYNGBYA PHORMIDIUM</i> , Var. <i>rivularis</i> ,	299
“ 22-26. <i>LYNGBYA PHORMIDIUM</i> , variable in diameter,	299
“ 27-29. <i>LYNGBYA ARENARIUM</i> , specimens from vicinity of Atlantic City, New Jersey,	299




EXPLANATIONS AND REFERENCES,

PLATE CCII.

Figures magnified 250 diameters.

		PAGE.
Figs. 1, 2.	LYNGBYA AESTUARII. forma <i>minus</i> , near New Jersey Sea Coast,	296
	“ “ Absecom, Perth Amboy, etc., compare Plate CC, fig. 11-16.	296
“ 3, 4.	LYNGBYA SUBTORULOSA, from Florida,	300
Fig. 5-8.	LYNGBYA RUPESTRE, specimens collected by Capt. J. D. Smith, on rocks, Falls of Deep Creek, Maryland,	300
“ 9-11.	LYNGBYA PAPYRINA, collected near New Haven, Conn.,	302
“ 12-14.	LYNGBYA INTERRUPTA, from Charlotte, Vermont,	302
“ 15-17.	LYNGBYA RETZII, from wet soil and wet rocks,	302
Figs. 18, 19.	LYNGBYA CATARACTA, water falls, Pennsylvania,	302
“ 20, 21.	LYNGBYA JULIANA, from Florida and Alabama,	301
“ 22, 23.	LYNGBYA TINCTORIA, from Arizona,	301
“ 24, 25.	LYNGBYA INUNDATA, margins of wayside ditches,	303
Fig. 26-31.	LYNGBYA PALLIDA, meadow grounds, New Jersey, Virginia, Florida, Pennsylvania,	298
Figs. 32, 33.	LEPTOTHRIX OCHRACEA, (not <i>Lyngbya ochracea</i>),	322









EXPLANATIONS AND REFERENCES,


PLATE CCIII.

Figures magnified 250 diameters.

	PAGE.
Figs. 1, 2. LEPTOTHRIX TENAX,	319
“ 3, 4. MICROCOLEUS HYALINUS,	307
Fig. 5-9. MICROCOLEUS ANGUIFORMIS, represents also <i>M. terrestris</i> , Var. <i>Vaucheria</i> ,	306, 305
Figs. 10, 11. MICROCOLEUS GRACILIS,	306
Fig. 12-14. MICROCOLEUS RAVENELII, from near Houston, Texas,	307













EXPLANATIONS AND REFERENCES,


PLATE CCIV.

Figures magnified 250 diameters.

	PAGE.
Figs. 1, 2. MICROCOLEUS (<i>Dasygloia</i>) AMORPHA, filaments in normal de-	
veloped condition,	304
Fig. 3-6. " young plants, spirally coiled, opening gradually,	305
" 7-9. " sheaths with one or more trichomes,	305
" 10-12. MICROCOLEUS PULVINATUS, sheaths with one, two or three tri-	
chomes,	305
Figs. 13, 14. " " two small thalli, less than natural	
size,	305*









EXPLANATIONS AND REFERENCES,

PLATE CCV.

Figures mostly magnified 250 diameters.

		PAGE.
Fig. 1.	MICROCOLEUS ANGUIFORMIS	306
Figs. 2, 3.	MICROCOLEUS HETEROTRICHUM, forma <i>Americana</i> , younger condition,	307
Fig. 4.	“ “ more advanced condition showing the two kinds of trichomes, sheath smooth,	308
“ 5.	“ “ older filament with sheath lamellate and fibrillose,	308
Figs. 6, 7.	ISACTIS FLUVIATILIS,	244, 245
Fig. 8.	SYMPLOCA LUCIFUGA, two vertical fascicles greatly magnified,	303
Figs. 9, 10.	“ “ natural size of plants,	303
“ 11, 12.	“ “ single trichomes in sheaths,	303
Fig. 13.	SYMPLOCA FRIESIANA, a caespitose cluster,	304
Figs. 14, 15.	“ “ single filaments from the cluster, with a single trichome,	304
“ 16, 17.	MICROCOLEUS TERRESTRIS, Var. <i>repens</i> , from moist earth,	305






EXPLANATIONS AND REFERENCES,

PLATE CCVI.

Figures magnified 250 diameters.

	PAGE
Fig. 1. OSCILLARIA SUBTILISSIMA,	309
" 2. OSCILLARIA TENERRIMA,	309
" 3. OSCILLARIA AMPHIBIA,	310
" 4. OSCILLARIA ELEGANS,	310
" 5. OSCILLARIA CRUENTA,	312
" 6. OSCILLARIA GRACILLIMA,	311
" 7. OSCILLARIA LEPTOTRICHA,	311
" 8. OSCILLARIA ANTLIARIA,	310
" 9. OSCILLARIA AERUGINEO-COERULEA,	311
" 10. OSCILLARIA VIOLACEA,	311
" 11-13. OSCILLARIA SUBFUSCA,	312
" 14. OSCILLARIA TENUIS,	313
" 15. OSCILLARIA CORTIANA,	313
" 16. OSCILLARIA LIMOSA,	313
" 17. OSCILLARIA CHALYBEA,	314
" 18. OSCILLARIA ANGUINA, (<i>subsalsa</i>),	214
" 19. OSCILLARIA NATANS,	314
" 20. OSCILLARIA NIGRA,	315
" 21. OSCILLARIA CHALYBEA,	314
Figs. 22, 23. OSCILLARIA PERCURSA,	315




EXPLANATIONS AND REFERENCES,

PLATE CCVII.

Figures magnified 250 diameters.

	PAGE.
Fig. 1-3. OSCILLARIA CRUENTA, parts of ordinary filaments,	312
“ 4-7 “ “ young, coiled filaments,	312
“ 8. OSCILLARIA BREVIS,	312
Figs. 9, 10. OSCILLARIA FROELICHI,	315
“ 11, 12. OSCILLARIA MAJOR,	316
“ 13, 15. OSCILLARIA GRATELOUPII,	316
“ 16, 17. OSCILLARIA BONNEMAISONII,	316
“ 18, 19. OSCILLARIA LITTORALIS,	317
“ 20, 22. OSCILLARIA PRINCEPS,	317
















EXPLANATIONS AND REFERENCES,

PLATE CCVIII.

Figures magnified 250 diameters.


	PAGE.
Figs 1, 2. OSCILLARIA IMPERATOR, smaller form,	317
“ 3, 4. “ “ typical form,	317
Fig. 5. LEPTOTHRIX HINNULEA,	320
Figs. 6, 7. BEGGIATOA LEPTOMITIFORMIS, two small mats; and single filaments,	319
Fig. 8-10. BEGGIATOA NIVEUM, two clusters, and single filaments more highly magnified,	318
Figs. 11, 12. LEPTOTHRIX CAESPITOSA, two caespitose clusters under low power,	320
Fig. 13. LEPTOTHRIX HERBACEA,	320
“ 14. LEPTOTHRIX OCHRACEA,	322
“ 15. LEPTOTHRIX RIGIDULA,	321
“ 16. LEPTOTHRIX TINCTORIA,	321
Figs. 17, 18. LEPTOTHRIX LAMINOSA,	321
Fig. 19. LEPTOTHRIX BULLOSA,	321
“ 20. “ “ single filaments under higher power,	321
“ 21. LEPTOTHRIX CALCICOLA, (<i>muralis</i>),	319




EXPLANATIONS AND REFERENCES,

PLATE CCIX.

	PAGE.
Fig. 1-3. CRENOTHRIX POLYSPORA, a section of filament with the cytoplasm divided longitudinally, and developed into spores; some of them developing clusters of filaments,	294
“ 4-9. Macrospores, or resting spores dividing and multiplying.	294
“ 10. Part of a filament under higher power showing the formation of macrospores,	294
Figs. 11, 12. Macrospores sliding out of the sheaths,	294
Fig. 13. Microgonidia gathered into masses, forming a mucus around themselves, <i>Palmella</i> -like,	294
“ 14. A small <i>Palmella</i> mass, gonidia developing into plants.	294
“ 15-17. Three sections of filaments under higher magnification, showing the division of the cytoplasm and formation of gonidia,	294
“ 18. A frequent appearance of a caespitose cluster,	294
“ 19. OSCILLARIA DETERSA	310
Figs. 20, 25. ASTEROTHRIX PERTYANA,	323
“ 22, 25. ASTEROTHRIX CREGINII,	322







EXPLANATIONS AND REFERENCES,

PLATE CCX.

		PAGE.
Fig. 1.	SPHIRILLUM UNDULA,	324
“ 2.	SPIRULINA JENNERI, two short sections,	323
“ 3.	SPIRULINA TENUISSIMA,	323
Figs. 4, 5.	SPIRULINA DUPLEX,	323
Fig. 6.	GLOEOTHECE CONFLUENS,	325
Figs. 7, 8.	APHANOTHECE PALLIDA,	325
“ 9, 10.	APHANOTHECE PRASINA,	325
Fig. 11.	SYNECHOCOCCUS AERUGINOSUS,	326
“ 12, 13, 15.	MERISMOPIDIA GLAUCA,	326
Fig. 14.	MERISMOPIDIA CONVOLUTA,	326
“ 16.	COELOSPHAERIUM KUETZINGIANUM,	327
Figs. 17, 18.	CLATHROCYSTIS AERUGINOSA, two perforated thalli, greatly enlarged,	327
Fig. 19.	“ “ teguments quickly dissolve and cells float in mass,	327
“ 20.	GOMPHOSPHAERIA APONINA,	328
Figs. 21, 22.	“ “ young conditions,	328
Fig. 23.	MICROCYSTIS PROTOGENITA,	328
“ 24.	POLYCYSTIS ICHTHYOBLABE,	330
“ 25.	ANACYSTIS PULVEREUS,	329
“ 26-31.	GLOEOCAPSA MAGMA, etc.,	331
“ 32.	GLOEOCAPSA, compare Plate CXCIV, figs. 24-28,	274
“ 33-37.	APHANOCAPSA VIRESCENS,	333
Figs. 38, 39.	APHANOCAPSA GREVILLEI,	333
“ 40, 41.	CHIROCOCCUS TURGIDUS,	334
Fig. 42.	CHIROCOCCUS COHERENS,	335
“ 43.	CHIROCOCCUS RUFESCENS,	335

